

ISSN 2518-1467 (Online),  
ISSN 1991-3494 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

# Х А Б А Р Ш Ы С Ы

---

---

**ВЕСТНИК**

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН

**THE BULLETIN**

THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN

PUBLISHED SINCE 1944

5

SEPTEMBER – OCTOBER 2019

---

---

ALMATY, NAS RK

---

*NAS RK is pleased to announce that Bulletin of NAS RK scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of Bulletin of NAS RK in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential multidiscipline content to our community.*

*Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабаршысы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабаршысының Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді мультидисциплинарлы контентке адалдығымызды білдіреді.*

*НАН РК сообщает, что научный журнал «Вестник НАН РК» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Вестника НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному мультидисциплинарному контенту для нашего сообщества.*

Б а с р е д а к т о р ы

х. ғ. д., проф., ҚР ҰҒА академигі

**М. Ж. Жұрынов**

Р е д а к ц и я а л қ а с ы:

**Абиев Р.Ш.** проф. (Ресей)  
**Абишев М.Е.** проф., корр.-мүшесі (Қазақстан)  
**Аврамов К.В.** проф. (Украина)  
**Аппель Юрген** проф. (Германия)  
**Баймуқанов Д.А.** проф., корр.-мүшесі (Қазақстан)  
**Байтулин И.О.** проф., академик (Қазақстан)  
**Банас Иозеф** проф. (Польша)  
**Берсимбаев Р.И.** проф., академик (Қазақстан)  
**Велесько С.** проф. (Германия)  
**Велихов Е.П.** проф., РҒА академигі (Ресей)  
**Гашимзаде Ф.** проф., академик (Әзірбайжан)  
**Гончарук В.В.** проф., академик (Украина)  
**Давлетов А.Е.** проф., корр.-мүшесі (Қазақстан)  
**Джрбашян Р.Т.** проф., академик (Армения)  
**Қалимолдаев М.Н.** проф., академик (Қазақстан), бас ред. орынбасары  
**Лаверов Н.П.** проф., академик РАН (Россия)  
**Лупашку Ф.** проф., корр.-мүшесі (Молдова)  
**Мохд Хасан Селамат** проф. (Малайзия)  
**Мырхалықов Ж.У.** проф., академик (Қазақстан)  
**Новак Изабелла** проф. (Польша)  
**Огарь Н.П.** проф., корр.-мүшесі (Қазақстан)  
**Полещук О.Х.** проф. (Ресей)  
**Поняев А.И.** проф. (Ресей)  
**Сагиян А.С.** проф., академик (Армения)  
**Сатубалдин С.С.** проф., академик (Қазақстан)  
**Таткеева Г.Г.** проф., корр.-мүшесі (Қазақстан)  
**Умбетаев И.** проф., академик (Қазақстан)  
**Хрипунов Г.С.** проф. (Украина)  
**Юлдашбаев Ю.А.** проф., РҒА корр.-мүшесі (Ресей)  
**Якубова М.М.** проф., академик (Тәжікстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабаршысы».

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде  
01.06.2006 ж. берілген №5551-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,  
<http://www.bulletin-science.kz/index.php/en/>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р  
д. х. н., проф. академик НАН РК  
**М. Ж. Журинов**

Р е д а к ц и о н н а я к о л л е г и я:

**Абиев Р.Ш.** проф. (Россия)  
**Абишев М.Е.** проф., член-корр. (Казахстан)  
**Аврамов К.В.** проф. (Украина)  
**Апель Юрген** проф. (Германия)  
**Баймуканов Д.А.** проф., чл.-корр. (Казахстан)  
**Байтулин И.О.** проф., академик (Казахстан)  
**Банас Иозеф** проф. (Польша)  
**Берсимбаев Р.И.** проф., академик (Казахстан)  
**Велесько С.** проф. (Германия)  
**Велихов Е.П.** проф., академик РАН (Россия)  
**Гашимзаде Ф.** проф., академик (Азербайджан)  
**Гончарук В.В.** проф., академик (Украина)  
**Давлетов А.Е.** проф., чл.-корр. (Казахстан)  
**Джрбашян Р.Т.** проф., академик (Армения)  
**Калимолдаев М.Н.** академик (Казахстан), зам. гл. ред.  
**Лаверов Н.П.** проф., академик РАН (Россия)  
**Лунашку Ф.** проф., чл.-корр. (Молдова)  
**Моход Хасан Селамат** проф. (Малайзия)  
**Мырхалыков Ж.У.** проф., академик (Казахстан)  
**Новак Изабелла** проф. (Польша)  
**Огарь Н.П.** проф., чл.-корр. (Казахстан)  
**Полещук О.Х.** проф. (Россия)  
**Поняев А.И.** проф. (Россия)  
**Сагиян А.С.** проф., академик (Армения)  
**Сатубалдин С.С.** проф., академик (Казахстан)  
**Таткеева Г.Г.** проф., чл.-корр. (Казахстан)  
**Умбетаев И.** проф., академик (Казахстан)  
**Хрипунов Г.С.** проф. (Украина)  
**Юлдашбаев Ю.А.** проф., член-корр. РАН (Россия)  
**Якубова М.М.** проф., академик (Таджикистан)

**«Вестник Национальной академии наук Республики Казахстан».**

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5551-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 2000 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18.

www: nauka-nanrk.kz, bulletin-science.kz

---

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

**M. Zh. Zhurinov**

E d i t o r i a l b o a r d:

**Abiyev R.Sh.** prof. (Russia)  
**Abishev M.Ye.** prof., corr. member. (Kazakhstan)  
**Avramov K.V.** prof. (Ukraine)  
**Appel Jurgen,** prof. (Germany)  
**Baimukanov D.A.** prof., corr. member. (Kazakhstan)  
**Baitullin I.O.** prof., academician (Kazakhstan)  
**Joseph Banas,** prof. (Poland)  
**Bersimbayev R.I.** prof., academician (Kazakhstan)  
**Velesco S.,** prof. (Germany)  
**Velikhov Ye.P.** prof., academician of RAS (Russia)  
**Gashimzade F.** prof., academician ( Azerbaijan)  
**Goncharuk V.V.** prof., academician (Ukraine)  
**Davletov A.Ye.** prof., corr. member. (Kazakhstan)  
**Dzhrbashian R.T.** prof., academician (Armenia)  
**Kalimoldayev M.N.** prof., academician (Kazakhstan), deputy editor in chief  
**Laverov N.P.** prof., academician of RAS (Russia)  
**Lupashku F.** prof., corr. member. (Moldova)  
**Mohd Hassan Selamat,** prof. (Malaysia)  
**Myrkhalykov Zh.U.** prof., academician (Kazakhstan)  
**Nowak Isabella,** prof. (Poland)  
**Ogar N.P.** prof., corr. member. (Kazakhstan)  
**Poleshchuk O.Kh.** prof. (Russia)  
**Ponyaev A.I.** prof. (Russia)  
**Sagiyani A.S.** prof., academician (Armenia)  
**Satubaldin S.S.** prof., academician (Kazakhstan)  
**Tatkeyeva G.G.** prof., corr. member. (Kazakhstan)  
**Umbetayev I.** prof., academician (Kazakhstan)  
**Khripunov G.S.** prof. (Ukraine)  
**Yuldashbayev Y.A.,** prof. corresponding member of RAS (Russia)  
**Yakubova M.M.** prof., academician (Tadjikistan)

**Bulletin of the National Academy of Sciences of the Republic of Kazakhstan.**

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5551-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://nauka-nanrk.kz/>, <http://bulletin-science.kz>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

A. S. Mukhamejanova<sup>1</sup>, G. A. Shalakhmetova<sup>2</sup>, A. N. Antipov<sup>3</sup>, Z. Alikulov<sup>1</sup>

<sup>1</sup>L. N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan,

<sup>2</sup>Al-Farabi Kazakh National University, Almaty, Kazakhstan,

<sup>3</sup>A. N. Bakh Institute of Biochemistry, Federal State Institution «Federal Research Centre  
«Fundamentals of Biotechnology» of the Russian Academy of Sciences, Moscow, Russia.

E-mail: akmaral1411@gmail.com; shalakhmetova@mail.ru; alex131313@yandex.ru; zer-kaz@mail.ru

## OPTIMIZATION OF CONDITIONS FOR THE ACTIVATION OF XANTHINE OXIDASE AND THE FORMATION OF NITRIC OXIDE IN FRESH SHEEP MILK

**Abstract.** For the detection of sheep milk XO activity a various natural and artificial antioxidants were examined. Among the natural antioxidants L-cysteine was more effective in the stabilization of XO in heated milk. XO of sheep milk activated by heat treatment in the presence of cysteine and molybdenum became able to convert nitrate and nitrite to nitric oxide (NO). Therefore, L-cysteine was used for double purposes: as the protector of enzyme active center against the oxidation during heat treatment of milk and as a reagent for S-nitrosothiol formation. Hypoxanthine, a natural substrate of XO, was the effective electron donor for NaR and NiR activities. Heat treatment of the milk in the presence of exogenous lecithin increased the activity of NaR and NiR of XO and CysNO formation. Thus, during the heat treatment: a) excess of exogenous phospholipids disintegrate the structure of MFGM and b) enzyme molecules denatured partially and their active center became available for exogenous cysteine, molybdenum, hypoxanthine and nitrate or nitrite.

**Key words:** milk, xanthine oxidase, NO, NO<sub>3</sub>, NO<sub>2</sub>, antioxidants, molybdenum, tungsten, phospholipids.

**Introduction.** At present, the pollution of environment with nitrates is a big problem in the world. In the environment and in human and animal body a microorganisms may convert nitrates to nitrites.

It is well known that nitrates (NO<sub>3</sub><sup>-</sup>) and nitrites (NO<sub>2</sub><sup>-</sup>) cause various diseases, including cancer. It is well known that nitrites irreversible bind to haemoglobin forming methaemoglobin which losses the ability to transport oxygen [1]. Deficiency of oxygen causes asphyxiation and it is particularly hazardous to health for babies. Furthermore, nitrites easily bind to primary amines, such as cadaverine, putrescine, spermidine and form potential carcinogens – nitrosamines [2, 3].

In 1980 one of the authors of this article for the first time observed that purified and homogeneous xanthine oxidase (XO) of cow's milk has the ability to reduce NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup> [5]. Later, other groups of scientists have found that animal XO converts NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup> to physiologically important gas – nitric oxide (NO) [6, 7, 8]. It is generally recognized that NO is one of the major biological messenger molecules, regulating blood pressure and blood flow, neurotransmission and brain function, immune system function, wound healing inhibition of platelet aggregation. NO is also involved in defense mechanisms against pathogens and some kinds of cancer cells [9]. In 1992 the nitric oxide was recognized as a molecule of year and in 1998 scientists studying its properties have been awarded the Nobel Prize [10].

Because of the ability to form nitric oxide the XO of milk harbors an antimicrobial activity. It is known that NO as the oxidant is a strong antibacterial agent. Antibacterial functions of XO are associated with peroxynitrite (ONOO<sup>-</sup>) which is the product of the reaction between NO and ·O<sub>2</sub> (superoxide anion). XO is also involved in protective an antiviral responses by catalyzing the conversion of retinaldehyde to retinoic acid. Retinoic acid derivatives can inhibit viral replication and, thus, preventing the spread of viral disease [11, 12].

Xanthine oxidoreductase or dehydrogenase/oxidase (XO; EC 1.1.3.22) – molybdenum and iron-containing flavoprotein. Each of the two subunits of XO contains one molybdenum center, two iron-sulfur center and one FAD. XO of the breast milk of women shows a molecular mass of 160 kDa subunit, XO of the cow's milk - 150 kDa. It is believed that the main biological function of XO is the catalyses the final step of purine oxidation in eucaryotes, it catalyzes the sequence of hydroxylations that convert hypoxanthine to xanthine, then to uric acid [13]. However, the enzyme has broad substrate specificity and is capable of reducing oxygen to generate the reactive oxygen species (ROS), superoxide and hydrogen peroxide, as well as oxidative transformation of pteridines and some aliphatic and aromatic aldehydes. Xanthine oxidase (XO) is not strongly specific to the oxidation of hypoxanthine or xanthine, it may catalyze the oxidation of about thirty nitrogen containing heterocycles and aldehydes [13]. Therefore because of its multifunctional enzymatic reactions XO is considered as a potential enzyme detoxifying different xenobiotics [14]. It is known that numerous heterocyclic xenobiotics (including pesticides) are carcinogens. Thus, in the case of contamination of milk with harmful xenobiotics the active XO makes possible their biotransformation into harmless forms. XO reduces nitrite (and nitrate), yielding reactive nitrogen species (RNS), such as nitric oxide and peroxynitrite [15].

In cow's milk XO localized in the inner membrane layer of the fat globules (XO content is about 8% of total proteins of cow milk fat globules) [16]. However, XO of milk fat globule membranes (MFGM) contains tenth times less molybdenum or does not contain it which results in a very low enzyme activity [17-19]. Study of the cofactor composition of XO isolated from cow's and women's milk showed that more than 85-90 % of enzyme molecules don't contain molybdenum in its active center and are inactive molecules [18]. The occurrence of xanthine oxidase in animal milk was a matter of slight controversy for a long time, because in the milk xanthine oxidation never been observed, although the levels of XO in the milk are relatively high. Presumably, this is due to the fact that before inclusion of Mo atoms in the active center of the enzyme, Mo-free molecules of XO already is associated with inner membrane of milk fat globules. As a result, of XO molecules become molecules without molybdenum. It is proposed that XO is involved in secretion of milk fat globules in a process dependent on the enzyme protein rather than on its enzymic activity [15].

Thus, in the presence of molybdenum atom in the active centre XO is active but in the absence of the metal it becomes inactive. Like other molybdoenzymes, in the active center XO contains so called molybdocofactor which has a pterin nature (therefore, it is called also molybdopterin) [20]. Molybdenum atom is bound to the pterin via its vicinal thiol groups (figure 1). This pterin is synthesized *in vivo* independently on the presence of molybdenum, i.e. in the absence of molybdenum cellular XO contains a normal amount of the cofactor. The cofactor is buried deeply within the interior of enzyme molecule and a tunnel-like structure makes it accessible to the appropriate substrates [21].




Figure 1 – Structure of molybdocofactor and its bond with molybdenum atom via vicinal SH-groups in the active center of XO [20]

**Materials and Methods.** *Preparation of the milk to the detection of the various enzymatic activities.* Before treatment in the milk of domestic animals added 10  $\mu$ M ethylenediaminetetraacetic acid (EDTA) to bind heavy metals. For boiling fresh milk is poured into the narrow conical tubes in a volume of 2 ml. Then, for further determine of the enzymatic activity the tubes are placed in water bath with 35  $^{\circ}$ C temperature and kept for 10 min, then using the special reagents different activities of XO are determined.

*Method for determination of the intrinsic activity of xanthine oxidase (XO).* To determine the intrinsic activity of XO 200  $\mu$ l aliquot of milk was mixed with 700  $\mu$ l 0.1 M sodium phosphate buffer containing

10  $\mu\text{M}$  EDTA, 5  $\mu\text{l}$  phenylmethylsulfonylfluor (for inhibition of protease activity). To this mixture was added 100  $\mu\text{M}$  10 mM hypoxanthine. The mixture was incubated at 30°C for 10 min under aerobic conditions. The proteins in the mixture is precipitated with trichloroacetic acid. After centrifugation, the amount of uric acid (under the influence of XO converted hypoxanthine to uric acid) in the supernatant was determined by measuring the absorbance of the reaction mixture in a spectrophotometer at 295 nm [22]. The amount of uric acid also is determined by its chemical oxidation with potassium permanganate ( $\text{K}_2\text{MnO}_4$ ) or hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), thereby forming allantoin [23]. Allantoin in biological liquids is easy to very accurately determined by a known method [24].

*Method for the determination of nitrate- and nitrite-reducing activity of XO.* Nitrate-reducing activity of XO determined by the disappearance of the added nitrate ( $\text{NO}_3^-$ ) or by the appearance of nitrite ( $\text{NO}_2^-$ ) in the reaction medium [5]. Nitrite-reducing activity of XO is determined by the disappearance of the nitrite to the reaction medium or by the appearance of nitric oxide (NO) [25].

**Results and Discussion.** *Proposed mechanisms of milk XO activation.* As mentioned above, milk XO in exists in molybdenum-free form and it is localized in the inner layer of MFGM. Therefore, the activation of milk XO requires the incorporation of exogenous molybdenum into its active center [4]. It is generally known that XO belongs to heat-stable enzymes – it remain active at 75-80°C temperature in several minutes [5]. However, at this temperature, the enzyme molecules undergo partial reversible denaturation. Therefore, one of possible ways for the availability of XO molecules for exogenous molybdenum is the disintegration of milk fat globule membranes and partial denaturation of enzyme molecules. Thus, during the heat treatment: a) excess of exogenous phospholipids disintegrate the structure of MFGM and b) enzyme molecules denatured partially and their active center became available for exogenous cysteine, molybdenum, hypoxanthine and nitrate or nitrite. Cysteine not only protects SH-groups of Mo-co against oxidation but also promotes the binding of molybdenum to the cofactor.

*Effects of different antioxidants in the stabilizing of heat-denatured milk XO.* Our preliminary experiments showed that the optimal heating temperature and its duration for detection of all activities in fresh milk XO were 80°C and 7 min (data not shown). Besides, the presence of 0.1 mM concentrations of cysteine and exogenous molybdenum was also necessary for the detection. Thus, to detect the activity of MFGM-bound enzyme is really required heat treatment and the addition of cysteine as an antioxidant and molybdenum to fresh sheep milk. In the next experiments we examined various natural and artificial antioxidants for the detection of milk XO activity. An natural antioxidants – cysteine, glutathione, lipoic acid and ascorbic acid, and artificial antioxidants – dithiotreitol, unithiol and mercaptoethanol (2,3-dimercapto-1-propanesulfonic acid) were used in different concentrations (table 1).

Results present in the table 1 show that among the natural antioxidants L-cysteine was more effective in the stabilization of XO in heat treated milk. In the same concentrations the effectiveness of cysteine containing antioxidant – glutathione was considerably lower. Apparently, this is due to the fact that tripeptide molecule of glutathione contains only one cysteine residue. Although both the oxidized (disulfide) and reduced (dihydro) forms of lipoic acid (dithiol) show antioxidant properties [26], its stabilizing effect for milk XO was lower than glutathione. Among the artificial antioxidants monothiol mercaptoethanol was more effective in the stabilization of heated milk XO. Although dithiotreitol and unithiol contain two sulfhydryl groups they don't stabilize heat treated milk XO. Recently we found that dithiotreitol in the presence of sulfanilamide forms stable blue complex with molybdate ion (on the basis of this finding we developed new fast method for molybdenum determination in the biological materials). Therefore, dithiotreitol associated with molybdenum does not show a stabilizing or antioxidant effect. The absence of such a stabilizing effect of unithiol remains unclear.

*Effect of different electron donors on associated activities of fresh sheep milk.* In the experiment for comparison various electron donors were used for own XO, NaR and NiR activities of fresh milk enzyme. Fresh sheep milk heated at 80°C in 7 min in the presence of 10  $\mu\text{M}$  EDTA, 0.1 mM cysteine, 0.1 mM  $\text{Na}_2\text{MoO}_4$  (or  $\text{Na}_2\text{WO}_4$ ). After cooling to the milk aliquots (200  $\mu\text{l}$ ) were added different electron donors separately (their concentrations are shown in the Table) and incubated for 15 min at 35°C. The final volume of reaction mixture was brought to 0.5 ml with 0.1 M Na-phosphate buffer (pH 6.0). After incubation the volume of the mixture was adjusted to 3 ml. The mixture was precipitated with acetic acid at pH 3.5 and centrifuged at 15000g in 15 min. In the supernatant the associated activities of milk XO were determined. In fresh untreated sheep milk none of associated activities was detected.



Table 1 – Effects of different antioxidants on the stability of XO during the heat treatment of fresh sheep milk in the presence of Na<sub>2</sub>MoO<sub>4</sub>

| Antioxidant     | Concentrations, mM | Associated activities |             |            |
|-----------------|--------------------|-----------------------|-------------|------------|
|                 |                    | XO                    | NaR         | NiR        |
| Cysteine        | 0.05               | 9.5 ± 1.4             | 75.2 ± 8.4  | 42.1 ± 6.3 |
|                 | 0.1                | 13.5 ± 1.7            | 93.5 ± 10.2 | 53.2 ± 6.8 |
|                 | 0.5                | 11.5 ± 2.4            | 78.7 ± 8.5  | 39.7 ± 7.1 |
| Glutathione     | 0.05               | 4.8 ± 0.9             | 66.3 ± 1.7  | 35.2 ± 4.2 |
|                 | 0.1                | 10.3 ± 1.6            | 78.6 ± 3.9  | 48.6 ± 8.7 |
|                 | 0.5                | 12.6 ± 2.1            | 94.3 ± 9.7  | 54.3 ± 8.2 |
| Ascorbic acid   | 0.05               | 7.6 ± 1.0             | 57.7 ± 9.7  | 42.7 ± 7.2 |
|                 | 0.1                | 9.5 ± 0.9             | 78.4 ± 4.9  | 48.6 ± 5.2 |
|                 | 0.5                | 10.3 ± 1.6            | 84.2 ± 9.3  | 50.2 ± 7.2 |
| Lipoic acid     | 0.05               | 7.3 ± 1.6             | 69.7 ± 9.2  | 40.4 ± 6.2 |
|                 | 0.1                | 9.5 ± 1.3             | 77.5 ± 15.0 | 51.2 ± 6.4 |
|                 | 0.5                | 8.9 ± 1.2             | 74.3 ± 9.9  | 48.2 ± 8.7 |
| Dithiotreitol   | 0.05               | 0                     | 0           | 0          |
|                 | 0.1                | 0                     | 0           | 0          |
|                 | 0.5                | 0                     | 0           | 0          |
| Unithiol        | 0.05               | 0                     | 0           | 0          |
|                 | 0.1                | 0                     | 0           | 0          |
|                 | 0.5                | 0                     | 0           | 0          |
| Mercaptoethanol | 0.05               | 9.2 ± 1.6             | 73.8 ± 9.2  | 41.3 ± 5.9 |
|                 | 0.1                | 10.7 ± 1.7            | 89.7 ± 11.3 | 53.0 ± 6.2 |
|                 | 0.5                | 9.2 ± 1.3             | 71.3 ± 9.7  | 40.4 ± 8.3 |

XO: nmoles of uric acid/0.1 ml/min; NaR: nmoles of NO<sub>2</sub>- formed/0.1 ml/min; NiR: nmoles of NO<sub>2</sub>- reduced/0.1 ml/min.

Table 2 – Effects of different electron donors on nitrate- and nitrite-reducing activity and own activities of sheep milk XO in the presence of cysteine

| Electron donors               | Associated activities |              |             |
|-------------------------------|-----------------------|--------------|-------------|
|                               | XO                    | NaR          | NiR         |
| 10 μ NADPH                    | 0.0                   | 0.0          | 0.0         |
| 10 μ NADH                     | 18.3 ± 2.7            | 114.7 ± 18.7 | 68 ± 9.4    |
| 10 μ FADH <sub>2</sub>        | 2.4 ± 0.6             | 21.3 ± 1.6   | 27.2 ± 4.2  |
| 0.1 mM hypoxanthine           | –                     | 92.8 ± 8.5   | 57.6 ± 7.6  |
| *Reduced methylviologen (MVH) | –                     | 128.6 ± 21.3 | 98.6 ± 14.3 |

XO: nmoles of uric acid/0.1 ml/min; NaR: nmoles of NO<sub>2</sub>- formed/0.1 ml/min; NiR: nmoles of NO<sub>2</sub>- reduced/0.1 ml/min; MVH: Reference 4.

Results in the table 2 show that the highest levels of the associated activities of milk XO observed with artificial electron donor – methylviologen reduced by dithionite (Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>). Among the physiological electron donors the highest associated activities were obtained using reduced NADH. Natural substrate of XO – hypoxanthine was also effective in the donating of electrons for NaR and NiR activities of the enzyme in fresh sheep milk. Thus, the electron transfer may also occur from the hypoxanthine (the reducing substrate) to nitrite and nitrate (the oxidizing substrates). Because in the presence of tungstate in the reaction mixture the associated activities were not detected, during nitrate reduction by hypoxanthine, the

XO molybdenum centre may participate directly in both the oxidative (hypoxanthine oxidation) and the reductive (nitrate and nitrite reduction) half-reactions.

*Detection of the products of nitrate- and nitrite reduction by XO in heat treated sheep fresh milk.* As mentioned above, it is found that animal XO converts  $\text{NO}_3^-$  and  $\text{NO}_2^-$  to physiologically important gas – nitric oxide (NO) [6-8]. NO effectively reacts with L-cysteine or reduced glutathione (GSH) at pH 7.0 and 7.4, to form orange-pink products of S-nitrosocysteine (CysNO) or S-nitrosogluthathione (GSNO). These products exhibited a peak absorbances at around 340 and 540 nm [25]. It is known that NADH is one of potential physiological electron donors for XO and it also has absorbance at 340 nm (reduced NADH exhibits strong UV absorption at 340 nm whilst the oxidized form has virtually no absorption at this wavelength). Therefore, to avoid mutual interference between the optical density of NADH and CysNO at 340 nm, instead of NADH we used hypoxanthine as an electron donor for NaR and NiR activities of milk XO (see above).

It was demonstrated that heat treatment (80°C, 10 min) of homogenic XO resulted in the release of molybdenum cofactor (Mo-co) from the active center of denatured enzyme molecule. During the heat treatment of XO ascorbic acid was the potential protector against the oxidation of released Mo-co. However, in the absence of ascorbic acid it quickly inactivated by oxygen (even in anerobic condions) [4]. Later we showed that glutathione and cysteine were the more powerful protectors for isolated Mo-co [27].

However ascorbic acid decomposes S-nitrosocysteine [28] and, therefore, we used L-cysteine as the protector against the oxidation of the cofactor in the active center of XO localized in MFGM. Thus, in our experiments L-cysteine was used for double purposes: as the protector of enzyme Mo-co against the oxidation during heat treatment of milk and as a reagent for S-nitrosothiol formation. For construction of calibration curve we used nitroprusside as a donor of NO. Increasing concentrations of nitroprusside from 10 nM to 1.0  $\mu\text{M}$  mixed with constant 0.1 mM concentration of cysteine in milk serum. Micromolar concentrations of nitroprusside release nanomolar concentrations of NO [20]. The serum was obtained by heating fresh milk, precipitating its proteins with acetic acid and centrifuging. In order to obtain a transparent milk serum, its filtration was carried out through an “Merk Millipore” membrane (33 mm Millex Filter Units).

For NO determination sheep milk mixed in the ratio of 1:1 with 0.2 M chlorinated phosphate buffer (PBS), pH 6.5, containing 10mM NEM, 2.5 mM EDTA [28], 0.2 mM  $\text{Na}_2\text{MoO}_4$  or  $\text{Na}_2\text{WO}_4$  and 0.1 mM cysteine. After incubation in 15 minutes at a temperature of 36°C the milk proteins were precipitated by diluted acetic acid added to the reaction mixture until pH 4.0. After centrifugation in the supernatant absorbance at 340 nm and 540 nm were measured.

For many years tungsten was considered to be a biological antagonist of molybdenum and was used for study of the properties and functions of molybdenum in Mo-enzymes. This was due to the fact that tungsten is able to replace molybdenum in Mo-enzymes, forming catalytically inactive analogs [30]. Therefore, to make sure that it is the molybdenum enzyme that catalyzes the formation of NO, instead of molybdenum we incubated the milk in the presence of tungsten (table 3).

Thus, the results obtained (table 3) convincingly show that the heat treatment of fresh sheep milk in the presence of exogenous molybdenum actually activates XO and the enzyme becomes capable of converting nitrate and nitrite to nitric oxide. However, when nitrates were used as a substrate NO formation was very low. At the same time, using nitrites as substrate resulted in 10 times higher amount of formed NO (i.e. CysNO) in comparison with nitrate substrates. The levels of CysNO determined by absorbances at 340 and 540 nm were completely different. It is likely that this was due to the difference in the sensitivity of the absorption at the ultraviolet and visible wavelengths of the spectrophotometer.

*Effect of exogenous phospholipid on the associated activities of sheep milk XO.* Phospholipids (PLs) are formed by glycerol, phosphoric acid, fatty acids and a hydroxy compound (e.g., choline, ethanolamine, serine, inositol). It is well known that milk fat globule membranes (MFGM) are composed of four main species of PLs: phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylinositol (PI) and phosphatidylserine (PS). The most abundant PLs in milk fat, expressed as percentage of total, are PE (26.4 % – 72.3 % of total PLs), PC (8.0 % – 45.5 %), PI (1.4 % – 14.1 %) and PS (2.0 % – 16.1 %). Of the lipids, phosphatidylcholine is largely located on the outside while the neutral lipids, phosphatidylethanolamine, phosphatidylserine and phosphatidylinositol are concentrated on the inner surface [31]. Amphiphilic properties of PLs derive from the presence of both a hydrophobic tail and a hydrophilic head

Table 3 – Formation of CysNO in fresh sheep milk after heat treatment in the presence of cysteine and molybdenum or tungsten

| Treatments of the milk  | Substrate in reaction mixture | Absorbance at: | Amount of CysNO, *nanomoles |
|---|-------------------------------|----------------|-----------------------------|
| Control – PBS only without milk   | NO <sub>3</sub> <sup>-</sup>  | 340 nm         | 0.0                         |
|   |                               | 540 nm         | 0.0                         |
|   | NO <sub>2</sub> <sup>-</sup>  | 340 nm         | 0.0                         |
|   |                               | 540 nm         | 0.0                         |
| Control – milk without heat treatment   | NO <sub>3</sub> <sup>-</sup>  | 340 nm         | 2.7 ± 0.3                   |
|   |                               | 540 nm         | 0.0                         |
|   | NO <sub>2</sub> <sup>-</sup>  | 340 nm         | 5.8 ± 0.2                   |
|   |                               | 540 nm         | 0.0                         |
| Heating + Na <sub>2</sub> MoO <sub>4</sub> at 80°C, 7 minutes                         | NO <sub>3</sub> <sup>-</sup>  | 340 nm         | 12.6 ± 0.8                  |
|   |                               | 540 nm         | 3.2 ± 0.2                   |
|   | NO <sub>2</sub> <sup>-</sup>  | 340 nm         | 35.5 ± 3.5                  |
|   |                               | 540 nm         | 7.3 ± 1.3                   |
| Heating + Na <sub>2</sub> WO <sub>4</sub> at 80°C, 7 minutes                          | NO <sub>3</sub> <sup>-</sup>  | 340 nm         | 0.0                         |
|   |                               | 540 nm         | 0.0                         |
|   | NO <sub>2</sub> <sup>-</sup>  | 340 nm         | 0.0                         |
|   |                               | 540 nm         | 0.0                         |
| Heating of milk without MoO <sub>4</sub> <sup>=</sup> or WO <sub>4</sub> <sup>=</sup> | NO <sub>3</sub> <sup>-</sup>  | 340 nm         | 0.0                         |
|   |                               | 540 nm         | 0.0                         |
|   | NO <sub>2</sub> <sup>-</sup>  | 340 nm         | 0.0                         |
|   |                               | 540 nm         | 0.0                         |
| *CysNO nanomoles/0.1 ml/min.  |                               |                |                             |

make them natural detergents. Thus, under high temperature phospholipids may effectively disrupt hydrophobic bonds between fatty acids, e.g., the integrity of all membrane layers of MFGM. Therefore, the destruction of MFGM makes the enzyme molecules available for exogenous molybdenum and its substrates (hypoxanthine, nitrate and nitrite).

In the next experiments, we examined the effect of the lecithin on NaR and NiR activities of heat-treated fresh sheep milk. Egg yolk lecithin from AppliChem (Panreac, ITW Companies, Darmstadt, Germany) was used in the experiments. The main phospholipids of this lecithin are phosphatidylcholine and phosphatidylethanolamine, i.e. they are components of the outer and inner layers of the membrane. The lecithin was dissolved in isopropyl alcohol and then added to the milk before heat treatment. The other components required for the heat treatment of milk have already been written above. Hypoxanthine was used an electron donor for NaR and NiR activities.

Table 4 – Effect of increasing concentrations of the lecithin on nitrate- and nitrite-reducing activities and CysNO formation in sheep milk

| Lecithin content                          | Associated activities |              | Amount of CysNO* |              |
|---|-----------------------|--------------|------------------|--------------|
|   | NaR                   | NiR          | From NaR         | From NiR     |
| Control                                   | 92.8 ± 8.5            | 57.6 ± 7.6   | 13.2 ± 1.2       | 33.7 ± 5.2   |
| 0.01 mg/ml                                | 91.2 ± 10.2           | 58.4 ± 6.8   | 14.3 ± 2.6       | 33.5 ± 7.3   |
| 0.05 mg/ml                                | 92.7 ± 9.5            | 57.8 ± 9.3   | 14.4 ± 1.8       | 34.7 ± 7.1   |
| 0.1 mg/ml                                 | 95.8 ± 11.6           | 62.9 ± 9.7   | 18.4 ± 3.2       | 42.9 ± 8.2   |
| 0.5 mg/ml                                 | 122.5 ± 14.7          | 117.6 ± 12.8 | 33.6 ± 4.5       | 78.9 ± 12.2  |
| 1.0 mg/ml                                 | 128.6 ± 12.8          | 135.2 ± 18.6 | 36.7 ± 7.3       | 93.8 ± 11.70 |
| *Results obtained from 340 nm absorbance. |                       |              |                  |              |

The results presented in table 4 show that exogenous lecithin in relatively high concentrations (0.5–1.0 mg/ml) increases the activity of NaR, NiR and the formation of CysNO. Apparently, during heat treatment, exogenous lecithin does partially decompose the membrane of MFGM of milk and enzyme molecules become more accessible. Thus, the results obtained may be important in the cleaning of animal milk contaminated with nitrates or nitrites. Thus, our results suggest the possible use of XO activation by heat treatment to remove nitrates from milk. Usually fresh sheep's milk is consumed after heat treatment.

**А. С. Мухамеджанова<sup>1</sup>, Г. А. Шалахметова<sup>2</sup>, А. Н. Антипов<sup>3</sup>, З. Аликулов<sup>1</sup>**

<sup>1</sup>Л. Н. Гумилев атындағы, Еуразия ұлттық университеті, Нур-Султан, Қазақстан,  
<sup>2</sup>Әл-Фараби атындағы Қазақ ұлттық университеті, Алматы, Қазақстан,  
<sup>3</sup>А. Н. Бах атындағы Биохимия институты, Ресей ғылым академиясы, Мәскеу, Ресей

### **ҚОЙДЫҢ САУМАЛ СҮТІНДЕГІ КСАНТИНОКСИДАЗАНЫ АКТИВТЕНДІРУ ЖӘНЕ АЗОТ ТОТЫҒЫНЫҢ ТҮЗІЛУ ЖАҒДАЙЛАРЫН ОҢТАЙЛАНДЫРУ**

**Аннотация.** Қойдың сүтіндегі КО-ның активтігін анықтау үшін әртүрлі табиғи және жасанды антиоксиданттар тексерілді. Табиғи антиоксиданттардың арасында L-цистеин қыздырылған сүттегі КО-ны күштірек тұрақтандыратын болып шықты. Цистеин мен молибденнің қатысуымен қыздырылған кезде активтенген сүттің КО-сы нитрат пен нитритті азот тотығына (NO) айналдыратын қабілетке ие болды. Сондықтан, цистеин екі мақсатта: қыздыру кезінде ферменттің активті орталығын тотығудан қорғайтын протектор және S-нитрозотиолдардың түзілуіне қажет реагент ретінде пайдаланылды. КО-ның табиғи субстраты – гипоксантин NaP мен NiP-дің активтіктері үшін күшті электрондар доноры бола алатыны анықталды. Сүтті сырттан берілген лецитиннің қатысуымен қыздырғанда NaP мен NiP-дің активтігі және CysNO-ның түзілуі жоғарылады. Сонымен, қыздыру кезінде: а) сырттан берілген фосфолипидтер СМГМ-ның құрылысын бұзды және б) ферменттің молекулалары жартылай денатурацияланады және оның нәтижесінде олардың активтік орталықтарына сырттан берілген цистеин, молибден, гипоксантин, нитрат немес нитрит ене алады.

**Түйін сөздер:** сүт, ксантиноксидаза, NO, NO<sub>3</sub>, NO<sub>2</sub>, антиоксиданттар,, молибден, вольфрам, фосфолипидтар.

**А. С. Мухамеджанова<sup>1</sup>, Г. А. Шалахметова<sup>2</sup>, А. Н. Антипов<sup>3</sup>, З. Аликулов<sup>1</sup>**

<sup>1</sup>Евразийский национальный университет им. Л. Н. Гумилева, Нур-Султан, Казахстан,  
<sup>2</sup>Казахский национальный университет им. аль-Фараби, Алматы, Казахстан,  
<sup>3</sup>Институт биохимии им. А. Н. Баха, Москва, Россия

### **ОПТИМИЗАЦИЯ УСЛОВИЙ ДЛЯ АКТИВАЦИИ КСАНТИНОКСИДАЗЫ И ОБРАЗОВАНИЯ ОКСИДА АЗОТА В ОВЕЧЬЕМ МОЛОКЕ**

**Аннотация.** Для обнаружения активности ксантиноксидазы (КО) в овечьем молоке были исследованы различные природные и искусственные антиоксиданты. Среди природных антиоксидантов L-цистеин был более эффективным в стабилизации КО прогревом молока. КО молока, активированная прогреванием в присутствии цистеина и молибдена, обладала способностью превращать нитрат и нитрит в оксид азота (NO). Поэтому, цистеин был использован для двух целей: в качестве протектора активного центра фермента от окисления во время термообработки и в качестве реагента для образования S-нитрозотиолов. Природный субстрат КО – гипоксантин был эффективным донором электронов для NaP и NiP активностей. Прогревание молока в присутствии экзогенного лецитина повышало NaP и NiP активности фермента и образование CysNO. Таким образом, во время термообработки: а) экзогенные фосфолипиды разрушают структуру МЖГМ и б) молекулы фермента частично денатурируются и в результате их активные центры становятся доступными для экзогенных цистеина, молибдена, гипоксантина и нитрата или нитрита.

**Ключевые слова:** молоко, ксантиноксидаза, NO, NO<sub>3</sub>, NO<sub>2</sub>, антиоксиданты,, молибден, вольфрам, фосфолипиды.

**Information about authors:**

Mukhamejanova A. S., PhD student, L. N. Gumilyov Eurasian National University, Astana, Kazakhstan; [akmaral1411@gmail.com](mailto:akmaral1411@gmail.com); <https://orcid.org/0000-0003-3238-1269>

Shalakhmetova G. A., candidate of biological sciences, Department of Molecular Biology and Genetics, Faculty of Biology and Biotechnology, Kazakh National University, al-Farabi, Almaty, Kazakhstan; [shalakhmetova@mail.ru](mailto:shalakhmetova@mail.ru); <https://orcid.org/0000-0002-8723-6010>

Antipov A. N., candidate of biological sciences, Federal State Institution «Federal Research Centre «Fundamentals of Biotechnology» of the Russian Academy of Sciences, Bakh Institute of Biochemistry, Moscow, Russia; [alex131313@yandex.ru](mailto:alex131313@yandex.ru); <https://orcid.org/0000-0002-8894-4190>

Alikulov Z., candidate of biological sciences, professor of Department of biotechnology and microbiology, L. N. Gumilyov Eurasian National University, Astana, Kazakhstan; [zer-kaz@mail.ru](mailto:zer-kaz@mail.ru); <https://orcid.org/0000-0002-0990-8918>

**REFERENCES**

[1] Reynolds J.D., Ahearn G.S., Angelo M., Zhang J., Cobb F., Stamler J.S. S-nitrosohemoglobin deficiency: A mechanism for loss of physiological activity in banked blood // *Proc. National Acad. Sciences*. 2007. N 104(43). P. 17058-17062.

[2] Bryan N.S. Nitrite in Nitric Oxide Biology: Cause or Consequence? A Systems-Based Review // *Free Radical Biology & Medicine*. 2006. N 41. P. 691-701.

[3] Gladwin M.T., Schechter A.N., Kim-Shapiro D.B., Patel R.P., Hogg N., Shiva S., Cannon R.O., Kelm M., Wink D.A., Espey M.G., Oldfield E.H., Pluta R.M., Freeman B.A., Lancaster Jr.J.R., Feelisch M., Lundberg J.O. The Emerging Biology of the Nitrite Anion // *Nature Chemical Biology*. 2005. N 1(6). P. 308-314.

[4] Mukhamejanova A.S., Kulataeva M.S., Shalakhmetova G.A., Alikulov Z. The effect of in vivo exogenous molybdenum on the activity of xanthine oxidase in milk and liver of a sheep // *News of NAS RK. Series of biology and medicine*. 2018. N 5(323). P. 253-241. ISSN 2518-1629 (Online), ISSN 2224-5308 (Print).

[5] Alikulov Z.A., L'vov N.P., Kretovich V.L. Nitrate and nitrite reductase activity of milk xanthine oxidase // *Biokhimiia*. 1980. N 45. P. 1714-1718.

[6] Millar T.M., Stevens C.R., Benjamin N., Eisenthal R., Harrison R., Blake D.R. Xanthine oxidoreductase catalyzes the reduction of nitrates and nitrite to nitric oxide under hypoxic conditions // *FEBS Letters*. 1998. N 427. P. 225-228.

[7] Zhang Z., Nauthon D., Winyard P.G., Benjamin N. Generation of nitric oxide by a nitrite reductase activity of xanthine oxidase: a potential pathway for nitric oxide formation in the absence of nitric oxide synthase activity // *Bioch. Biophys. Res. Comm.* 1998. N 249. P. 767-772.

[8] Hord N.G, Ghannam J.S, Garg H.K, Berens P.D., Bryan N.S. Nitrate and nitrite content of human, formula, bovine, and soy milks: implications for dietary nitrite and nitrate recommendations // *Breastfeed Med*. 2011. N 6(6). P. 393-399.

[9] Andrew M., Garg H., Couglin J., Bryan N. Nutritional epidemiology in the context of nitric oxide biology: Risk-Benefit evaluation for dietary nitrite and nitrate // *Nitric Oxide*. 2010. N 22. P. 110-119.

[10] Bryan N.S., Bian K., Murad F. Discovery of the nitric oxide signaling pathway and targets for drug development // *Frontiers in Bioscience*. 2009. P. 1-18.

[11] Taibi G., Paganini A., Gueli M.C., Ampola F., Nicotra C.M. Xanthine oxidase catalyzes the synthesis of retinoic acid // *J Enzyme Inhib.* 2001. N 16(3). P. 275-85.

[12] Taibi G., Nicotra M., Concetta A. Xanthine oxidase catalyzes the oxidation of retinol // *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2007. N 22:4. P. 471-476.

[13] Harrison R. Physiological Roles of Xanthine Oxidoreductase // *Drug Metabolism Reviews*. 2004. Vol. 36, P. 363-375. doi:10.1081/DMR-120037569

[14] Beedham C. Molybdenum hydroxylases. In: «Enzyme systems that metabolize drug and xenobiotics» Ed. Costas Ioannidis // John Wiley & Sons Ltd. 2001. P. 146-188.

[15] Harrison R. Milk xanthine oxidase: Properties and physiological roles. // *International Dairy Journal*. 2006. Vol. 16, Iss. 6. P. 546-554.

[16] Mondy B.L., Keenan T.W. Butyrophilin and xanthine oxidase occur in constant molar proportions in milk lipid globule membrane but vary in amount with breed and stage of lactation // *Protoplasma*. 1993. N 177. P. 32-36.

[17] Godber B., Sanders S., Harrison R., Eisenthal R., Bray R. More or 95% of xanthine oxidase in human milk is present as the demolybdo form, lacking molybdopterin // *Biochem. Soc. Trans.* N 25. P. 519-529.

[18] Bray R.S, Lowe D., Godber B., Harrison R., Eisenthal R. Properties of xanthine oxidase from human milk: grossly deficient in molybdenum and substantially deficient in iron-sulfur centers // *Agency for Scientific Publications*. 1999. P. 775-778.

[19] Atmani D., Benboubetra M., Harrison R. Goat's milk xanthine oxidoreductase is grossly deficient in molybdenum // *J. Dairy Res*. 2004. N 71. P. 7-13.

- [20] Kramer P. Steven, Jean L. Johnson, Anthony A. Ribeiroll, David S. Millingtonll, K. V. Rajagopalan. The Structure of the Molybdenum Cofactor // *The Journal of Biological Chemistry*. 1987. Vol. 262, N. 34. P. 16357-16363.
- [21] Kisker C., Schindelin H., Rees D.C. Molybdenum-cofactor-containing enzymes: structure and mechanism // *Annual Review of Biochemistry*. 1997. N 66. P. 233-267.
- [22] Suzuki Go, Okamoto Ken, Kusano Teruo, Yoko Matsuda, Akira Fuse, Hiroyuki Yokota. Evaluation of Neuronal Protective Effects of Xanthine Oxidoreductase Inhibitors on Severe Whole-brain Ischemia in Mouse Model and Analysis of Xanthine Oxidoreductase Activity in the Mouse Brain // *Neurol Med Chir*. 2015. N 55(1). P. 77-85.
- [23] Mirko P. The Mechanism for the Conversion of Uric Acid into Allantoin and Dehydro-Allantoin // *A New Look at an old Problem. Tetrahedron*. 1986. N 42. P. 747-751.
- [24] Vogels G., Van der Drift C. Differential analyses of glyoxylate derivatives. // *Analytical Biochemistry*. 1970. N 33. P. 143-157.
- [25] Kuo W.N., Kocis J.M., Nibbs J. Nitrosation of cysteine and reduced glutathione by nitrite at physiological pH // *Front Biosci*. 2003. N 8. P. 62-69.
- [26] Rochette L., Ghibu S., Richard C., Zeller M., Cottin Y., Vergely C. Direct and indirect antioxidant properties of  $\alpha$ -lipoic acid and therapeutic potential // *Mol Nutr Food Res*. 2013. N 57(1). P. 114-125.
- [27] Alikulov Z.A., Mendel R.R. Molybdenum cofactor from tobacco cell cultures and milk xanthine oxidase: involvement of sulfhydryl groups in dimerization activity of cofactor // *Biochemie und Physiologie der Pflanzen*. 1984. N 179(8). P. 693-705.
- [28] Stsiapura V., Stepuro I., Morozkina T., Lewis S., Smith L. S-Nitrosoglutathione formation at gastric pH is augmented by ascorbic acid and by the antioxidant vitamin complex // *Journal Pharmaceutical Biology*. 2018. Vol. 56.
- [29] Marley R. Formation of nanomolar concentrations of S-nitroso-albumin in human plasma by nitric oxide // *Free Radical Biology & Medicine*. 2001. N 31(5). P. 688-696.
- [30] Kletzin A., Adams M. W. Tungsten in biological systems // *FEMS Microbiol Rev*. 1996. N 18(1). P. 5-63.
- [31] Garcia C., Lutz N.W., Confort-Gouny S., Cozzone P.J., Armand M., Bernard M. Phospholipid fingerprints of milk from different mammals determined by 31P NMR: Towards specific interest in human health // *Food Chem*. 2012. N 135. P. 1777-1783.

---

---

**Publication Ethics and Publication Malpractice  
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

**ISSN 2518-1467 (Online), ISSN 1991-3494 (Print)**

<http://www.bulletin-science.kz/index.php/en/>

Редакторы *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*  
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 10.10.2019.  
Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
13,9 п.л. Тираж 500. Заказ 5.