

**ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ГЫЛЫМ АКАДЕМИЯСЫНЫҢ

Д.В.Сокольский атындағы «Жанармай,
катализ және электрохимия институты» АҚ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
АО «Институт топлива, катализа и
электрохимии им. Д.В. Сокольского»

N E W S

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
JSC «D.V. Sokolsky institute of fuel, catalysis
and electrochemistry»

**SERIES
CHEMISTRY AND TECHNOLOGY**

2 (434)

MARCH - APRIL 2019

PUBLISHED SINCE JANUARY 1947

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of chemistry and technologies scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Химия және технология сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруды. Web of Science зерттеушілер, авторлар, баспашилар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енүі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия химии и технологий» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Бас редакторы
х.ғ.д., проф., ҚР ҮҒА академигі М.Ж. Жұрынов

Редакция алқасы:

Агабеков В.Е. проф., академик (Белорус)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Ресей)
Газалиев А.М. проф., академик (Қазақстан)
Ергожин Е.Е. проф., академик (Қазақстан)
Жармағамбетова А.К. проф. (Қазақстан), бас ред. орынбасары
Жоробекова Ш.Ж. проф., академик (Қырғыстан)
Иткулова Ш.С. проф. (Қазақстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Қазақстан)
Баешов А.Б. проф., академик (Қазақстан)
Бұркітбаев М.М. проф., академик (Қазақстан)
Джусипбеков У.Ж. проф. корр.-мүшесі (Қазақстан)
Молдахметов М.З. проф., академик (Қазақстан)
Мансуров З.А. проф. (Қазақстан)
Наурызбаев М.К. проф. (Қазақстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф. академик (Қазақстан)
Стрельцов Е. проф. (Белорус)
Тәшімов Л.Т. проф., академик (Қазақстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Тәжікстан)
Фарзалиев В. проф., академик (Әзірбайжан)

«ҚР ҮҒА Хабарлары. Химия және технология сериясы».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №1089-Ж мерзімдік басылым тіркеуіне қойылу туралы күзелік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://chemistry-technology.kz/index.php/en/archiv>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д.х.н., проф.,академик НАН РК **М. Ж. Журинов**

Р е д а к ц и о н на я к ол л е г и я:

Агабеков В.Е. проф., академик (Беларусь)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Россия)
Газалиев А.М. проф., академик (Казахстан)
Ергожин Е.Е. проф., академик (Казахстан)
Жармагамбетова А.К. проф. (Казахстан), зам. гл. ред.
Жоробекова Ш.Ж. проф., академик (Кыргызстан)
Иткулова Ш.С. проф. (Казахстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Казахстан)
Баешов А.Б. проф., академик (Казахстан)
Буркитбаев М.М. проф., академик (Казахстан)
Джусипбеков У.Ж. проф. чл.-корр. (Казахстан)
Мулдахметов М.З. проф., академик (Казахстан)
Мансуров З.А. проф. (Казахстан)
Наурызбаев М.К. проф. (Казахстан)
Рудик В. проф.,академик (Молдова)
Рахимов К.Д. проф. академик (Казахстан)
Стрельцов Е. проф. (Беларусь)
Ташимов Л.Т. проф., академик (Казахстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Таджикистан)
Фарзалиев В. проф., академик (Азербайджан)

«Известия НАН РК. Серия химии и технологии».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,

<http://chemistry-technology.kz/index.php/en/arhiv>

© Национальная академия наук Республики Казахстан, 2019

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142,
Институт органического катализа и электрохимии им. Д. В. Сокольского,
каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief
doctor of chemistry, professor, academician of NAS RK **M.Zh. Zhurinov**

Editorial board:

Agabekov V.Ye. prof., academician (Belarus)
Volkov S.V. prof., academician (Ukraine)
Vorotyntsev M.A. prof., academician (Russia)
Gazaliyev A.M. prof., academician (Kazakhstan)
Yergozhin Ye.Ye. prof., academician (Kazakhstan)
Zharmagambetova A.K. prof. (Kazakhstan), deputy editor in chief
Zhorobekova Sh.Zh. prof., academician (Kyrgyzstan)
Itkulova Sh.S. prof. (Kazakhstan)
Mantashyan A.A. prof., academician (Armenia)
Praliyev K.D. prof., academician (Kazakhstan)
Bayeshov A.B. prof., academician (Kazakhstan)
Burkitbayev M.M. prof., academician (Kazakhstan)
Dzhusipbekov U.Zh. prof., corr. member (Kazakhstan)
Muldakhmetov M.Z. prof., academician (Kazakhstan)
Mansurov Z.A. prof. (Kazakhstan)
Nauryzbayev M.K. prof. (Kazakhstan)
Rudik V. prof., academician (Moldova)
Rakhimov K.D. prof., academician (Kazakhstan)
Streltsov Ye. prof. (Belarus)
Tashimov L.T. prof., academician (Kazakhstan)
Toderash I. prof., academician (Moldova)
Khalikov D.Kh. prof., academician (Tadzhikistan)
Farzaliyev V. prof., academician (Azerbaijan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of chemistry and technology.
ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)
The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10893-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Editorial address: Institute of Organic Catalysis and Electrochemistry named after D. V. Sokolsky
142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22,
e-mail: orgcat@nursat.kz

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

N E W S

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES CHEMISTRY AND TECHNOLOGY

ISSN 2224-5286

<https://doi.org/10.32014/2019.2518-1491.19>

Volume 2, Number 434 (2019), 55 – 60

UDC 547.99**A.I. Zhussupova, Y.S. Ikhsanov, A.A. Mamutova, G.E. Zhusupova**

Al-Farabi Kazakh National University

Center for Physical and Chemical Methods of Analysis

Republic of Kazakhstan, 050012, Almaty, Karasay Batyr St., 95a

Aizhan.Zhussupova@kaznu.kz, erbol.ih@gmail.com, Alua.Mamutova@kaznu.kz, zhusupova@gmail.com

**COMPARATIVE ANALYSIS OF THE NONPOLAR FRACTION
OF THE AERIAL AND UNDERGROUND PARTS OF THE *LIMONIUM*
GMELINII PLANTS BY THE GC-MS METHOD**

Abstract. This article discusses the chemical composition of the non-polar fraction obtained from the aerial and underground parts of *Limonium gmelinii* (Willd.) harvested in the Almaty region in 2018. Extracts were obtained by the method of solvent extraction with hexane and studied by means of chromato-mass spectrometry on a gas chromatograph with a mass-selective detector.

As a result, 22 compounds were isolated from the aerial part of *Limonium gmelinii* plant, out of which dominating both quantitatively and qualitatively are hydrocarbons, such as genicosane and eicosane and tricosane; quinoline and phytol derivatives were isolated as well. High content of phytol acetate related to derivatives of acyclic diterpene alcohols has been revealed. In addition, the higher alcohol 3,7,11,15-tetramethyl-2-hexadecan-1-ol has been identified.

Analysis of the non-polar extract obtained from the roots of *Limonium gmelinii* allowed us to isolate 14 compounds, out of which the major share in the amount belongs to the esters of higher carboxylic acids and hydrocarbons. Esters are represented as ethyl esters of hexadecanoic, oleic and linoleic acids with prevalence of polyene acids in them. Among the hydrocarbons, tridecane, tetradecane, heptadecane, octadecane and tetracosane are the most abundant.

Keywords: *Limonium gmelinii*, non-polar fraction, GC-MS, chemical composition.

Introduction

Limonium is a widespread plant genus with about 300 species present on the territory of Central and Western Asia. On the territory of CIS, 35 *Limonium* species have been identified. Usually it is accommodated on coniferous soils and dry mountain ranges, mainly in the south-eastern part of Europe, Caucasus, and Central Asia. In Kazakhstan, out of 18 species the most significant and productive is *Limonium gmelinii* (Willd.). The area of its distribution includes major part of Kazakhstan, Western and Eastern Siberia, European part of Russia, Central Asia, Southeastern Europe, Western China and Mongolia. Currently, the roots and the aerial parts of *L.gmelinii* plants have been introduced into the medicine and State Pharmacopoeia of the Republic of Kazakhstan, harmonized with the European Pharmacopoeia [1-8]; with effective drugs obtained and standardized on the basis of its roots [9-12].

The purpose of this work is a comparative analysis of the chemical composition of hexane extracts obtained from the aerial and underground parts of *L.gmelinii* plants, in order to establish their effect on the biological activity of the studied extracts.

Methods

Extracts from the aerial and underground parts of *L.gmelinii* plants were obtained by the method presented below:

A weighed amount of pre-treated, dried and crushed vegetable raw materials, according to their quality indicators, corresponding to the requirements of the State Pharmacopoeia of the Republic of Kazakhstan, weighing 10 g, was extracted with hexane at a ratio of 1:5 for 48 hours. The extract obtained was filtered and concentrated to 10 ml. The resulting total non-polar extract was studied on a gas chromatograph with a mass selective detector model Agilent 7890N/ 5973N (Agilent Technologies, USA). At the same time, the volume of the gas phase to be taken equals 1.00 μ l; the sample introduction temperature is 250°C without division of the flow. Separation was performed using a DB-35 MS chromatographic capillary column with a length of 30 m, an inner diameter of 0.25 mm and a film thickness of 0.25 μ m at a constant carrier gas (helium) rate of 1 ml/min. The temperature of chromatography was programmed from 40°C to 200°C with a heating rate of 10°C/min.

The software Agilent MSD ChemStation, version 1701EA) was used to control the gas chromatography system, record and process the obtained results and data. Data processing included determination of retention times, peak areas, as well as processing of spectral information obtained using a mass spectrometric detector. The Wiley 7th edition and NIST'02 libraries were used to decipher the mass spectra obtained (the total number of spectra in the libraries exceeds 550,000 items).

Results and discussion

As a result of studying the hexane fraction obtained from the aerial part of plants of the *Limonium gmelinii* species, 22 compounds were identified, most of which belong to the class of hydrocarbons (Table 1 and Figure 1).

Table 1 – Research data by GC-MS of the aerial part of *Limonium gmelinii* plant species

No.	Retention time, min	Name of compound	Content %
1	8.05	Dodecane	0.20
2	9.52	Tridecane	0.50
3	10.91	Tetradecane	1.21
4	11.54	Dodecane, 2,6,10-trimethyl-	0.60
5	12.23	Pentadecane	2.44
6	12.91	Dichloroacetic acid tridecyl ester	0.57
7	13.49	Hexadecane	3.50
8	13.88	1,2-dihydro-2,2,4-trimethyl Quinoline	0.72
9	14.48	2,6,10,14-tetramethyl-Pentadecane	1.16
10	14.68	Heptadecane	3.14
11	15.35	1-Decanol, 2-hexyl	0.36
12	15.,69	Tetracontan	1.29
13	15.81	Octadecane	2.71
14	16.45	Phytol acetate	12.88
15	16.99	Nonadecan	2.80
16	17.07	3,7,11,15-Tetramethyl-2-hexadecan-1-ol	5.17
17	17.20	2-Pentadecanone 6,10,14-trimethyl	2.46
18	18.48	Eicosane	1.93
19	20.47	Genicosan	7.82
20	22.53	1,2-Benzenedicarboxylic acid, butyl 2-ethylhexyl ester	0.35
21	22.87	Phytol	0.75
22	27.16	Tricosane	47.44
Total			100.00

As can be seen from the data presented in Table 1, the tricosane hydrocarbon (47.44%) dominates in the obtained hexane extract from the aerial part of the plants *Limonium gmelinii*. A high content of phytol acetate (12.88%) related to derivatives of acyclic diterpene alcohols has been established, in addition, the highest alcohol 3,7,11,15-tetramethyl-2-hexadecan-1-ol, detected in the amount of 5.17%, has been identified.




Figure 1 – GC-MS chromatogram of the aerial part of the plant species *Limonium gmelinii*

As can be seen from the data presented in Figure 1, the largest peaks are observed at 16.45 minutes and 27.16 minutes, which advises the extracts that are dominant in the extract, phytol acetate and tricosane, the other compounds are found in significant concentrations such as 3,7,11,15-tetramethyl-2-hexadecan-1-ol, genicosan, hexadecane are also clearly marked on the chart.

From the literature sources it is known that the underground parts of plants, as a rule, contain a relatively small amount of non-polar compounds, since the latter, phytol, in particular, are involved in the process of biosynthesis at the stage of photosynthesis and predominate in chloroplasts. The accumulation of other non-polar and weakly polar compounds, such as hydrocarbons, is more characteristic of the aerial parts of plants, while in the underground parts of plants, respectively, water-soluble carbohydrates and polyphenols dominate [13-23].

In the underground part of the studied plants, 14 compounds were identified (Table 2 and Figure 2).

Table 2 – Research data by GC-MS of the roots of plants of the *Limonium gmelinii* species

No.	Retention time, min	Name of compound	Content, %
1	6.53	Undecane	0.94
2	8.05	Dodecane	2.66
3	9.52	Tridecan	4.07
4	10.91	Tetradecane	4.15
5	10.99	7-Tetradecene, (Z)	0.89
6	12.23	9-methylheptadecane	3.22
7	13.49	Heptadecane	7.62
8	13.83	Phenol, 2,4-bis (1,1-dimethylethyl) -	1.72
9	13.88	Quinoline, 1,2-dihydro-2,2,4-trimethyl-	4.18
10	15.69	Tetracosan	3.43
11	15.81	Octadecane	5.26
12	19.78	Hexadecanoic acid ethyl ester	24.51
13	25.40	Oleic acid ethyl ester	11.16
14	25.78	Ethyl linoleic ester	26.19
Total			100.00

It was established on table 2 that the distinctive feature of the hexane extract of the roots of the studied plants is the predominance of esters of higher carboxylic acids in the amount of 61.86% and hydrocarbons – 32.24%.

The esters are represented as ethyl esters of hexadecanoic, oleic and linoleic acids, with the predominance of polyene acids (60.38%) in them, of which hexadecanoic acid ethyl ester and linoleic acid ethyl ester are the most abundant.




Figure 2 – GC-MS chromatogram of the roots of the plant species *Limonium gmelinii*

From the data presented in Figure 2, it can be seen that most of the dominant in the non-polar fraction of the roots of the plant *Limonium gmelinii* substances, namely Hexadecanoic acid ethyl ester, Oleic acid ethyl ester and Ethyl linoleic ester, belong to the class of organic esters is allocated at the final stages of analysis. The quinoline and phenol derivatives stand out between 1 and 5 minutes of the process. Most of the analysis recorded various compounds belonging to the class of hydrocarbons.

Conclusion

As a result, it was established that in the hexane extract obtained from the aerial part of the plant *Limonium gmelinii*, the dominant groups of substances are hydrocarbons among which the tridecane, tetradecane, heptadecane, octadecane and tetracosane dominate, 76.09%. Also in hexane extract obtained from the aerial part of the plant *Limonium gmelinii* was identified derivatives of phytol and quinoline.

The hexane extract obtained from the roots of the plant *Limonium gmelinii* is dominated by fatty acid esters whose total content is 61.86% in contrast to the aerial part of the plant. In addition, a significant amount of quinoline derivatives was found in the hexane extract from the roots of the plant *Limonium gmelinii*, while in the aerial part of the plant it is represented in trace amounts.

The obtained data can be used to improve the processing of the roots and the aerial parts of the plant *Limonium gmelinii* with the aim of obtaining drugs on its basis.

Funding

This work was carried out within the framework of the grant financing program of the SC MES RK AR05134034 “Development and creation of high-performance gels based on pharmacopoeial wild plants of Kazakhstan and their integrated research.”

УДК 547.99

А.И. Жусупова, Е.С. Ихсанов, А.А. Мамутова, Г.Е. Жусупова

Әл-Фараби атындағы Қазак ұлттық университеті

Физико-химиялық әдістер анализі орталығы

Қазақстан Республикасы, 050012, Алматы қ., Карасай батыра к., 95а,

**LIMONIUM GMELINII ӨСІМДІГІНІЦ ЖЕРҮСТІ ЖӘНЕ ЖЕРАСТЫ БӨЛІКТЕРІНІЦ
ПОЛЯРСЫЗ ФРАКЦИЯЛАРЫНЫЦ GC-MS ӘДІСІМЕН САЛЫСТАРМАЛЫ АНАЛИЗІ**

Аннотация. Бұл мақалада 2018 ж. Алматы облысында өндөлген *Limonium gmelinii* (Willd.) (кermек Гмелина) өсімдігінің жерасты және жерүсті бөліктерінен алынған полярсыз фракциялардың химиялық құрамы қарастырылады. Экстракттар гексанмен сұйық экстракциялау әдісімен алынып, масс-селективті детекторлы газ хроматографта хромато-масс спектрометрия арқылы зерттелді.

Нәтижесінде *Limonium gmelinii* өсімдігінің жерүсті бөлігінен 22 қосылыс, оның ішінде сапалық және сандық тұрғыдан көмірсүтектер гениказан, эйказан және триказан басым, сонымен қатар хинолин мен фитол туындылары бөлінді. Ациклді дитерпен спирттер туындыларының құрамы жоғары екендігі аныкталды, одан бөлек 3,7,11,15-тетраметил-2-гексадекан-1-ол жоғарғы спирті идентификацияланды.

Limonium gmelinii тамырынан алынған полярсыз экстракттар анализі 14 қосылыс бөліп алуға мүмкіндік берді, олардың негізгі сандық бөлігін жоғарғы карбон қышқылдарының эфилері мен көмірсүтектер құрайды. Құрделі эфирлер полиен қышқыларының басымдығымен линоль, олеин гексадекан қышқылдарының этил эфирлері күйінде ұсынылған. Көмірсүтектер ішінде көп кездесетіні тридекан, тетрадекан, гептадекан, октадекан мен тетракозан.

Түйін сөздер: *Limonium gmelinii*, полярсыз фракция, хромато-масс спектрометрия, химиялық құрам.

УДК 547.99

А.И. Жусупова, Е.С. Ихсанов, А.А. Мамутова, Г.Е. Жусупова

Казахский национальный университет имени аль-Фараби

Центр физико-химических методов анализа

Республика Казахстан, 050012, г.Алматы, ул.Карасай батыра, 95а,

**СРАВНИТЕЛЬНЫЙ АНАЛИЗ НЕПОЛЯРНОЙ ФРАКЦИИ НАДЗЕМНОЙ
И ПОДЗЕМНОЙ ЧАСТЕЙ РАСТЕНИЙ LIMONIUM GMELINII МЕТОДОМ GC-MS**

Аннотация. В данной статье рассматривается химический состав неполярной фракции, полученной из надземной и подземной частей растения *Limonium gmelinii* (Willd.) (кермек Гмелина), заготовленного в Алматинской области в 2018 году. Экстракти были получены методом жидкостной экстракции гексаном и изучены посредством хромато-масс спектрометрии на газовом хроматографе с масс-селективным детектором.

В результате исследования из надземной части растения *Limonium gmelinii* было выделено 22 соединения, доминирующими из которых, как в количественном, так и в качественном плане, являются углеводороды, такие как гениказан и эйказан и триказан; также были выделены производные хинолина и фитола. Установлено высокое содержание производных ациклических дитерпеновых спиртов, помимо этого идентифицирован высший спирт 3,7,11,15-тетраметил-2-гексадекан-1-ол.

Анализ неполярного экстракта, полученного из корней *Limonium gmelinii* позволил выделить 14 соединений; из них основную долю в количестве составляют эфиры высших карбоновых кислот и углеводороды. Сложные эфиры представлены в форме этиловых эфиров гексадекановой, олеиновой и линолевой кислот с преобладанием в них эфиров полиеновых кислот. Среди углеводородов в наибольшем количестве присутствуют тридекан, тетрадекан, гептадекан, октадекан и тетракозан.

Ключевые слова: *Limonium gmelinii*, неполярная фракция, хромато-масс спектрометрия, химический состав.

REFERENCES

- [1] Bajtenov MS (2001) Flora Kazahstana: rodovoj kompleks flory v 2-h t. Almaty: Gylym. ISBN 9965-07-036-9 (in Russ).

- [2] Kukenov MK (1999) Botanicheskoe resursovedenie Kazahstana. Almaty: Gylym. ISBN: 5-628-02318-3 (in Russ).

- [3] Sokolov PD (1984). Rastitel'nye resursy SSSR. L.: Nauka. ISBN 5-02-026723-6 (in Russ).
- [4] Komarov VL (1952) Flora SSSR. M.: AN SSSR. ISBN 978-5-458-42425-7 (in Russ).
- [5] Pavlov NI (1961) Flora Kazahstana – Alma-Ata: Nauka. T. VII. S. 79-80. ISBN 978-5-4458-5989-5 (in Russ).
- [6] Grudzinskaja LM, Gemedzhieva NG (2012) Spisok lekarstvennyh rastenij Kazahstana. Almaty. ISBN 978-601-80248-6-3 (in Russ).
- [7] Gosudarstvennaja farmakopeja Respubliki Kazahstan. T.1. (2008). Almaty: Izdatel'ckij dom "Zhibek zholy". ISBN 9965-759-97-98 (in Russ).
- [8] European Pharmacopoeia. 9th ed. (2018) Strasbourg. ISBN 978-3-7692-6815-7.
- [9] Korul'kina LM., Zhusupova GE., Shul'ts EE., Erzhanov KB. (2004), Fatty-acid composition of two Limonium plant species, Chemistry of natural compounds, 40: 417-419. DOI: 10.1007/s10600-005-0002-5.
- [10] Korul'kina LM, Shul'ts EE, Zhusupova GE, Abilov ZhA, Erzhanov KB, Chaudri MI (2004) Biologically active compounds from *L. gmelinii* and *L. popovii*. I, Chemistry of natural compounds, 40: 465-468. DOI: 10.1007/s10600-005-0012-3.
- [11] Zhusupova GE, Abil'kaeva SA (2006) Flavanes from *Limonium gmelinii*. II, Chemistry of natural compounds, 42: 112-113. DOI: 10.1007/s10600-006-0052-3.
- [12] Zhusupova GE (2009) Medical University Bulletin [Vestnik Medicinskogo Universiteta] 1: 105-112. (in Russian).
- [13] Jakushkina NI, Bahtenko EJ (2004) Fiziologija rastenij. Moskva. ISBN 5-691-01353-H.
- [14] Zeiger E, Moller IM, Murphy A, Taiz L (2018) Plant Physiology and Development. – 6th rev.ed. – Oxford University Press Inc. ISBN: 9781605357454.
- [15] Gadetskaya AV, Tarawneh AH, Zhusupova GE, Gemejyeva NG, Cantrell CL, Cutler SJ, Ross SA (2015) Sulfated phenolic compounds from *Limonium caspium*: Isolation, structural elucidation, and biological evaluation. Fitoterapia, 104, 80-85. DOI: 10.1016/j.fitote.2015.05.017.
- [16] Stankovic SM, Petrovi M, Godjevac D, Stevanovi ZD (2015) Screening inland halophytes from the central Balkan for their antioxidant activity in relation to total phenolic compounds and flavonoids: Are there any prospective medicinal plants? J arid env 120: 26-32. DOI: 10.1016/j.jaridenv.2015.04.008.
- [17] Rozentsvet OA, Nesterov VN, Bogdanova ES (2014) Membrane-forming lipids of wild halophytes growing under the conditions of Prieltonie of South Russia, Phytochem 105: 37-42. DOI: 10.1016/j.phytochem.2014.05.007.
- [18] Akhani H, Malekmohammadi M, Mahdavi P, Ghariyani A, Chase MW (2013) Phylogenetics of the Irano-Turanian taxa of *Limonium* (Plumbaginaceae) based on ITS nrDNA sequences and leaf anatomy provides evidence for species delimitation and relationships of lineages. Bot. J. Linn. Soc. 171: 519-550. DOI: 10.1111/bj.12015.
- [19] Gancedoa NN, Medeirosa D, Milaneze-Gutierrezb MA, Mello JP (2018) Morpho-anatomical characters of *Limonium brasiliense* leaves Revista Brasileira de Farmacognosia 28: 513-519 DOI: 10.1016/j.bjp.2018.05.014.
- [20] Lin LC, Chou CJ (2000) Flavonoids and phenolics from *Limonium sinense*. Planta Medica. 66: 382-383. DOI: 10.1055/s-2000-8547.
- [21] Komekbay Zh. N., Halmenova Z. B., Umbetova A. K., Bisenbay A.G (2018) Phytochemical analysis and development of production of biologically active complex on the basis of raw *Melissa officinalis* l. News of NAS RK. Series of chemistry and technology 427: 53 – 58 <https://doi.org/10.32014/2018.2518-1491>.
- [22] Umbetova A.K., Slan G.O., Omarova A.T., Burasheva G.Sh., Abidkulova K. T. (2018) The study of chemical composition of atraphaxis virgata from the almaty region. News of NAS RK. Series of chemistry and technology 428: 52 – 55 <https://doi.org/10.32014/2018.2518-1491>.
- [23] Utegenova L.A., Nurlybekova A.K., Aisa H., Jenis J. (2018) Liposoluble constituents of fritillaria pallidiflora. News of NAS RK. Series of chemistry and technology 432: 156 – 162 <https://doi.org/10.32014/2018.2518-1491.38>.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации
в журнале смотреть на сайте:

www:nauka-nanrk.kz

<http://chemistry-technology.kz/index.php/en/arhiv>

ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)

Редакторы: *М. С. Ахметова, Т. А. Апендиев, Аленов Д.С.*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 04.04.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
4 п.л. Тираж 300. Заказ 2.

*Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*