

ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Д.В.Сокольский атындағы «Жанармай,
катализ және электрохимия институты» АҚ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
АО «Институт топлива, катализа и
электрохимии им. Д.В. Сокольского»

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
JSC «D.V. Sokolsky institute of fuel, catalysis
and electrochemistry»

SERIES
CHEMISTRY AND TECHNOLOGY

3 (435)

MAY - JUNE 2019

PUBLISHED SINCE JANUARY 1947

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of chemistry and technologies scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Химия және технология сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия химии и технологий» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Б а с р е д а к т о р ы
х.ғ.д., проф., ҚР ҰҒА академигі **М.Ж. Жұрынов**

Р е д а к ц и я а л қ а с ы:

Ағабеков В.Е. проф., академик (Белорус)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Ресей)
Газалиев А.М. проф., академик (Қазақстан)
Ергожин Е.Е. проф., академик (Қазақстан)
Жармағамбетова А.К. проф. (Қазақстан), бас ред. орынбасары
Жоробекова Ш.Ж. проф., академик (Қырғыстан)
Иткулова Ш.С. проф. (Қазақстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Қазақстан)
Баешов А.Б. проф., академик (Қазақстан)
Бүркітбаев М.М. проф., академик (Қазақстан)
Джусипбеков У.Ж. проф. корр.-мүшесі (Қазақстан)
Молдахметов М.З. проф., академик (Қазақстан)
Мансуров З.А. проф. (Қазақстан)
Наурызбаев М.К. проф. (Қазақстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф. академик (Қазақстан)
Стрельцов Е. проф. (Белорус)
Тәшімов Л.Т. проф., академик (Қазақстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Тәжікстан)
Фарзалиев В. проф., академик (Әзірбайжан)

«ҚР ҰҒА Хабарлары. Химия және технология сериясы».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №1089-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://chemistry-technology.kz/index.php/en/arihiv>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д.х.н., проф., академик НАН РК **М. Ж. Журинов**

Р е д а к ц и о н н а я к о л л е г и я:

Агабеков В.Е. проф., академик (Беларусь)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Россия)
Газалиев А.М. проф., академик (Казахстан)
Ергожин Е.Е. проф., академик (Казахстан)
Жармагамбетова А.К. проф. (Казахстан), зам. гл. ред.
Жоробекова Ш.Ж. проф., академик (Кыргызстан)
Иткулова Ш.С. проф. (Казахстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Казахстан)
Баешов А.Б. проф., академик (Казахстан)
Буркитбаев М.М. проф., академик (Казахстан)
Джусипбеков У.Ж. проф. чл.-корр. (Казахстан)
Мулдахметов М.З. проф., академик (Казахстан)
Мансуров З.А. проф. (Казахстан)
Наурызбаев М.К. проф. (Казахстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф. академик (Казахстан)
Стрельцов Е. проф. (Беларусь)
Ташимов Л.Т. проф., академик (Казахстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Гаджикистан)
Фарзалиев В. проф., академик (Азербайджан)

«Известия НАН РК. Серия химии и технологии».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© Национальная академия наук Республики Казахстан, 2019

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142,
Институт органического катализа и электрохимии им. Д. В. Сокольского,
каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK **M.Zh. Zhurinov**

E d i t o r i a l b o a r d :

Agabekov V.Ye. prof., academician (Belarus)
Volkov S.V. prof., academician (Ukraine)
Vorotyntsev M.A. prof., academician (Russia)
Gazaliyev A.M. prof., academician (Kazakhstan)
Yergozhin Ye.Ye. prof., academician (Kazakhstan)
Zharmagambetova A.K. prof. (Kazakhstan), deputy editor in chief
Zhorobekova Sh.Zh. prof., academician (Kyrgyzstan)
Itkulova Sh.S. prof. (Kazakhstan)
Mantashyan A.A. prof., academician (Armenia)
Praliyev K.D. prof., academician (Kazakhstan)
Bayeshov A.B. prof., academician (Kazakhstan)
Burkitbayev M.M. prof., academician (Kazakhstan)
Dzhusipbekov U.Zh. prof., corr. member (Kazakhstan)
Muldakhmetov M.Z. prof., academician (Kazakhstan)
Mansurov Z.A. prof. (Kazakhstan)
Nauryzbayev M.K. prof. (Kazakhstan)
Rudik V. prof., academician (Moldova)
Rakhimov K.D. prof., academician (Kazakhstan)
Streltsov Ye. prof. (Belarus)
Tashimov L.T. prof., academician (Kazakhstan)
Toderash I. prof., academician (Moldova)
Khalikov D.Kh. prof., academician (Tadjikistan)
Farzaliyev V. prof., academician (Azerbaijan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of chemistry and technology.
ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10893-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://chemistry-technology.kz/index.php/en/arhiv>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Editorial address: Institute of Organic Catalysis and Electrochemistry named after D. V. Sokolsky
142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22,
e-mail: orgcat@nursat.kz

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES CHEMISTRY AND TECHNOLOGY

ISSN 2224-5286

<https://doi.org/10.32014/2019.2518-1491.26>

Volume 3, Number 435 (2019), 32 – 37

UDC 541.128

Zh.K. Kairbekov¹, A.S. Maloletnev², I.M. Jeldybayeva^{1,3}*, D.Z. Abilmazhinova³

¹SSE Research Institute of New Chemical Technologies and Materials, Almaty, Kazakhstan;

²Moscow Mining Institute NITU MISiS, Moscow, Russia;

³Kazakh State Women's Pedagogical University, Almaty, Kazakhstan;

abilmazhinova.1985@mail.ru

**DATA ON ANTIOXIDANT ACTIVITY OF HUMIC SUBSTANCES
OF LOW-MINERALIZED SLUDGE SULPHIDE MUD (PELOIDS)
AND THEIR CLASSIFICATION
(Review)**

Abstract. In this work, an overview of the antioxidant activity of humic substances of low-mineralized sludge sulphide muds (peloids). Though these substances show the greatest biological and therapeutic activity, in comparison with similar substances of other sources (keonardits, coals, etc.) they are insufficiently studied. Formed in the reduction conditions, molecules of humin substances in low-mineralized silt sulphidic mud possess a unique structure, and excludes a possibility of education in the course of receiving carcinogenic the alkilnic the halogenderivants and thereof urgent is a research in given the directions.

Key words: antioxidatic properties, activity, humin substances, peloida.

Peloid therapy or mud cure is one of the earliest methods of therapy using natural components, which have the biggest adaptogene potential. Being customary stimulants, they have an expressed curative influence [1,2].

The peloid therapy can be considered as a unique method for sanitation, since the range of deceases, on which it has positive influence, is quite wide [3]. Peloids have immunomodulatory effect, they cause natural adaptive reactions of human body [4,5], produce anti-inflammatory, desensitizing, antineoplastic [2,6,7], anaesthetic and absorbable effect, improve hemodynamics and lymphdynamics, reduce the activity of exudative and infiltrative processes, soften commissural structures [9]. The mud cure is applied when treating various pathologies: arthritis, respiratory illness, dermatological deceases, gastrointestinal problems, gynaecological deceases, trophic ulcers, hepatitis, inflammatory diseases of lungs, paradontium [10].

The therapeutic effects of mud are based on the combination of actions of closely related thermal, mechanical, chemical and biological factors. Depending on physical and chemical composition of the mud, the severity of the biological reactions of each of them is different. The literature sources describe studies devoted to the study of the peloid composition, but they all characterize mineral component of the mud, and almost do not include functional ability of the organic component. The insufficient knowledge of the organic substances of peloids caused their underestimation in the mechanism of effect on the body [11].

The peloids are divided into the following groups basing on the content of organic substances: peat, sapropelic, sludge sulphide [12]. A number of studies [10] has showed a high biological effect of sludge sulphide mud, which is associated with the components of the composition of the organic fraction. Ë

Sludge sulphide mud are dividing into the following categories basing on the content of sulphides: weakly sulphide (0.01–0.15 % of wet mud), medium sulphide (0.15–0.5 %), heavy sulphide (exceeding 0.5 %). Depending on the salinity of mud solution, low-mineralized (1.0–15.0 g/l), medium-mineralized (15.0–35 g/l), highly mineralized (35–150 g/l), saline-saturated (exceeding 150 g/l) sulphide muds [13] are distinguished.

The therapeutic significance of sludge sulphide mud is related to high viscous-plastic and thermal properties, content of large amounts of sulphides (exceeding 0.2 %) and organic substances (approximately 5 % on dry basis), including humic acids and their salts, lipids and hemicellulose, and presence of vitamins, enzymes and hormones [14]. Groups of nonspecific and specific compounds are distinguished from the composition of organic substances of peloids. Non-specific compounds include lipids, carbohydrates, amino acids, enzymes, keratinoid pigments, vitamin complexes, and others [10, 15-17].

A higher content of organic substances in peloids is represented by specific organic components – humic substances [16, 18, 19], which content is equal to 45–90 %.

Scientists of all countries are making attempts to isolate and study properties of organic components of peloids to use them independently as biologically active natural compounds and introduce them into public health practice. The study of fractions of humic substances of peloids: himatomelanic, humic, fulvic acids is a considerable interest for creating innovative pharmacotherapeutic drugs based on them, including those with high antioxidant activity [20-24].

Humic substances are a combination of biothermodynamically stable compounds that form, plant and animal residues having no analogues in living organisms in the process of decomposition and biotransformation, which are distinguished by dark color, polydispersity, high molecular weights [25-27]. All humic substances are formed as a result of the postmortal transformation of organic humification residues, process of transformation of decomposition products having different composition and origin of organic residues into humic substances [28].

The difference of this group of natural organic substances from others lies in their stoichiometric structure and stochastic nature related to the peculiarities of formation resulting from the selection of biothermodynamically stable structures. The variety of molecular forms of humic substances is determined by the mechanism of biochemical reactions that accompany formation and conditions of geochemical transformation [28, 29].

Classification of humic substances is based on various solubility of these substances in acids and alkalis. According to this classification, they are divided into the following fractions: fulvic, himatomelanic, humic, humus acids and humin-conglomerate of esters of acids and their organo-mineral sorption complexes with clay minerals, insoluble in most solvents [28-30].

Humic acids are insoluble at $\text{pH} < 2$, in a dissolved state they have a dark brown colour, in dry state – black powder or flakes [27]. They consist of aromatic core and peripheral part formed by specific chains.

The himatomelanic acids are an alcohol-soluble fraction, which have a cherry-red colour in solution and were first isolated in 1889 and described by the German biochemist, doctor Ernst Felix Hoppe-Zeiler [25].

Fulvic acids are soluble in the whole range of medium acidity, have a variable colour in the range from straw-yellow to orange colour. They vary from humic and himatomelanic acids by greater oxidation and lower carbon content, as well as greater hydrophilicity [27].

Humus acids are the sum of humic, himatomelanic, and fulvic acids. They have more folded, complex, stochastic structure. Presumably, being supramolecular structures, humus acids are destroyed during classic fractionation of humic substances. Besides, mechanical mixing of already obtained fractions following separation does not have the same set of properties, that humus acids have [14, 25].

Free radical activity is a fundamental property of humic substances [31, 32]. Studying the structure of various natural biopolymers, we can judge the reactivity of the specified substances and their possible transformations in one or another environment. The nature and properties of stable organic free radicals of humic substances has been investigated in a large number of works [32-33]. The content of aromatic structures in the composition of humic substances determines their low solubility in water and high stability during transformation. Electrophilic substitution reactions occur mainly at benzene nuclei, and side chain alkyl are primarily subjected to oxidation.

It is known that one of the most important characteristics of humic substances is high content of paramagnetic centers in them, which presence may be related to free radicals. Presence of paramagnetic centers increases, if the aromaticity of the humic series compounds increases [26]. Paramagnetism of humus acids and their fractions consists of at least three types of paramagnetic components: radicals with a predominant localization of the unpaired electron on the heteroatomic peripheral groups; high-molecular

systems with a sufficiently developed poly-conjugation chain with structural defects; as well as complexes of metals with variable valence [26].

The effect of humic substances on the intensity of biochemical processes is explained by their participation in reductive-oxidative reactions due to the presence of polyphenolic and quinoid groups. During the enzymatic oxidation of polyphenols, quinones are generated through seven quinones-type intermediate compounds [28].

In a similar manner, the electrons of reactive oxygen species and electrons in the molecules of humic substances are blocked.

The study of the antioxidant activity of humic substances obtained from various sources, today is an important task. The reductive-oxidative properties of soil humic substances [26], peat [34] under the influence of various chemical and physical and chemical factors have already been determined. At the same time, the study of the reductive-oxidative properties of humic substances of peloids is not well understood. Although these substances exhibit the greatest biological and therapeutic activity, compared with similar substances of other sources (keonardites, coals, etc.). Formed under reductive conditions, molecules of humic substances of low-mineralized sludge sulphide have a unique structure, and it excludes the possibility of generation when obtaining carcinogenic alkyl halogen derivatives. It means that the research in this field becomes important.

The methods for determining the antioxidant activity are based on the principles of direct or indirect measurement of the rate or completeness of reaction of antioxidants with appropriate reagents, among which amperometry and manometry have the greatest interest. For the quantitative determination of antioxidants, the amperometric method seems to be the most reliable, since it allows directly measuring the content of all antioxidants in a sample. It is based on the measurement of the electric current that is generated during the electrochemical oxidation of the investigated substance (or mixture of substances) on the surface of the work electrode at a certain potential. Under conditions of amperometric detection, compounds containing a hydroxyl group are well oxidized, the limit of their detection lies in the range of 10^{-9} – 10^{-12} g, and under favourable conditions, some compounds are determined at a level of 10^{-15} .

The antioxidant activity of humic acid solutions with various concentrations of 1 %, 0.1 %, 0.01 %, 0.001 % was measured in works [35-39] (Table 1). Quercetin was used as a reference material ($C_{15}H_{10}O_7$). To build an analytical curve of quercetin, recommended as a standard for this device, the All-Russia Scientific Research Institute of Metrological Service sequentially recorded signals of standard solutions of quercetin in the order of their concentrations increasing. According to the obtained results, curves of the peak area (signal value) dependence on quercetin concentration were built.

An arithmetic average of five measurements was taken as a result (relative standard deviation no more than 5 %), and a calibration curve was built (Figure 1).




Figure 1 - Dependence of the peak area on the concentrations of quercetin

Table 1 - The value of the total content of antioxidants.

Name	Concentration	Signal value (area), HAc	CCA, mg / ml (standard quercetin)
Humic acid	0.001%	516	Менее 0.0001
Humic acid	0.01%	3602	0.00058
Humic acid	0.1%	14571	0.00387
Humic acid	1%	44177	0.043

The value of the humic acid signal of 0.001 %, concentration – 516 nAc indicates that the total content of antioxidants is less than 0.0001 mg/ml. Increasing the concentration of humic acids, the value of digital signals, and total content of antioxidants increases. When the concentration of humic acids is 0.01 %, the total content of antioxidants is 0.00058 mg/ml, at a concentration of 0.1 % – 0.00387 mg/ml, and when the concentration of humic acids is equal to 1 %, the value of the total content of antioxidants reaches 0.043 mg/ml. The obtained data suggests that humic acids have antioxidant activity, which value increases, if the concentration is increased.

If several years ago humic substances were used mainly in agriculture and animal farming, today their use has proven to be in demand in pharmacy. The research of their physical, chemical and pharmacological properties will improve the effectiveness and accessibility of peloid therapy for patients, and will allow making dosed treatment. Peloid preparations are easier packed and transported, do not require special conditions for their storage. Waste material (mud) is disposed in the mud baths, but it still keep valuable substances. The active components isolated from these “wastes” can later be used to isolate humic substances and produce peloid drugs.

The Kazakhstan market of medicines is filled with foreign antioxidant drugs, and this affects their final price paid by the consumer. The country needs domestic drugs that can generate high competition. Therefore, the research related to the creation of innovative medicines based on humic substances of peloids has particular importance and relevance.

УДК 541.128

Ж.К. Қаирбеков¹, А.С. Малолетнев², И.М. Джелдыбаева^{1,3}*, Д.З. Абильмажинова³

¹ Жаңа химиялық технологиялар мен материалдар ҒЗИ, Алматы қаласы, Қазақстан

² Тау-кен институты, ҰЗТУ МҚжәнеБИ, Мәскеу

³ Қазақ мемлекеттік қыздар педагогикалық университеті, Алматы қаласы, Қазақстан

ЛАЙ СУЛЬФИДТІ БАТПАҚТАН АЛЫНҒАН ГУМИНДІК ЗАТТАРДЫҢ АНТИОКСИДАНТТЫ БЕЛСЕНДІЛІГІ МЕН ОЛАРДЫҢ КЛАССИФИКАЦИЯСЫ ТУРАЛЫ

(Шолу)

Аннотация. Бұл мақалада лай сульфидті батпақтан алынған (пелоид) гуминді заттардың антиоксиданттық белсенділігіне шолу жасалды. Бұл заттар басқа да көздерден алынған ұқсас заттармен салыстырғанда (кеонардит, көмір және т.б.) ішіндегі ең биологиялық және терапевтік белсенділікке ие болғанымен, олар аз зерттелген. Қалыптасу жағдайында туындаған лай сульфидті батпақтардағы гуминді заттардың молекулалары бірегей құрылымға ие және оларды алу барысында канцерогенді алкил галоген туындыларының пайда болу мүмкіндігін жоққа шығарады, нәтижесінде осы саладағы зерттеулердің маңыздылығы зор екендігі қалыптасады.

Түйін сөздер: антиоксиданттық белсенділік, гуминді заттар, пелоид.

УДК 541.128

Ж.К. Қаирбеков¹, А.С. Малолетнев², И.М. Джелдыбаева^{1,3}*, Д.З. Абильмажинова³

¹ ДГП НИИ Новых химических технологий и материалов, г. Алматы, Казахстан:

² Горный институт НИТУ МИСиС, Москва:

³ Казахский государственный женский педагогический университет г. Алматы, Казахстан

О АНТИОКСИДАНТНОЙ АКТИВНОСТИ ГУМИНОВЫХ ВЕЩЕСТВ НИЗКОМИНЕРАЛИЗОВАННЫХ ИЛОВЫХ СУЛЬФИДНЫХ ГРЯЗЕЙ (ПЕЛОИДОВ) И ИХ КЛАССИФИКАЦИЯ

(Обзор)

Аннотация. В данной работе сделан обзор об антиоксидантной активности гуминовых веществ низкоминерализованных иловых сульфидных грязей (пеллоидов). Хотя данные субстанции проявляют наибольшую биологическую и терапевтическую активность, по сравнению с аналогичными субстанциями

других источников (кеонардитов, углей и др.) они недостаточно изучены. Образованные в восстановительных условиях, молекулы гуминовых веществ низкоминерализованных иловых сульфидных грязей обладают уникальным строением, и исключает возможность образования в процессе получения канцерогенных алкильных галогенопроизводных, и вследствие этого актуальным становится исследование в данном направлении.

Ключевые слова: антиоксидантная активность, гуминовые вещества, пелоиды.

Information about authors:

Kairbekov Zhaksyntay Kairbekovich - Doctor of Chemical Sciences, Professor, Head of the Laboratory of SSE Scientific Research Institute New Chemical Technologies and Materials RSE al-Farabi Kazakh National University, Karasai batyr 95A, working tel.: 8 (727) 2921786, cells.: 87011584400, 87071384400, e-mail: zh_kairbekov@mail.ru, ORCID 0000-0002-0255-2330, <https://orcid.org/0000-0002-0255-2330>;

Anatoly Maloletnev - Doctor of Chemical Sciences, Professor, National Research Technological University "MISiS", Moscow, Russia, e-mail: anstanmal@mail.ru, ORCID 0000-0003-1952-9660, <https://orcid.org/0000-0003-1952-9660>;

Jeldybaeva Indira Mukhametkerimovna - PhD-doctor, Senior Lecturer Kazakh State Women's Pedagogical University, Aiteke-bi st. 99, working tel.: 8(727)2923732, mob.: 87079669796, 87026669796, e-mail: jeldybaeva@gmail.com, ORCID 0000-0002-1524-4046, <https://orcid.org/0000-0002-1524-4046>;

Abilmazhinova Didar Zamanbekovna - Ph.D doctoral student of the 2nd course of the Kazakh State Women's Pedagogical University, Aiteke-bi st. 99, mob. +77773878785, e-mail: abilmazhinova.1985@mail.ru, ORCID 0000-0001-7362-4963, <https://orcid.org/0000-0001-7362-4963>

REFERENCES

- [1] Dubishev A.V., Men'shich L.V. Minor Study of humic substances influence on peloids renal excretory function // Bulletin of Samara Scientific Center RAN. Samara, 2010, vol.12, №1 (8). P.2023-2026 (in Russ.).
- [2] Leshynski A.F, Zuza Z.I. Mud and pharmacotherapy in inflammatory diseases / Kiev, 1985.184 p. (in Russ.).
- [3] Kalenini S.V. Physiology of mud as a special case of non-specific adaptation of the organism // Vopr.kurortologii. 2003. №4. P.52-53 (in Russ.).
- [4] Mechuperenko O.N. Mud - natural biogenic stimulators, healing mechanisms of action / Provizor. 1998. №6. P.54-57(in Russ.).
- [5] Suramfer M.J. Smoking, estrogen and prevention of hear disease in women / Mayo elin, proc.-1989. Vol.84. P.1553-1557 (in Russ.).
- [6] Semionova M.A. Chemical and pharmaceutical, organizational and economic feasibility of application of humic acids of peloids: Avtoref.diss.kand.farm.nauk / Samara, 2006, 20 p.
- [7] Shustov L.P. Extracts of silt sulfide mud and the rationale for their use in clinical practice / Vopr.kurortologii. 1999. №6. P.35-37 (in Russ.).
- [8] Ulashchik V.S., Lukomsky I.V. Total fizioterapiya: textbook / Minsk, 2004. 327 p. (in Russ.)
- [9] Shustov L.P. Extracts of silt sulfide mud and healing of use, Tomsk, 1996. 182 p. (in Russ.)
- [10] Agaplov A.I., Mezhevalova N.I.,Avvakumova N.P. Peloidopreparaty humic series as a means of increasing the efficiency of ieloidoterapii under new conditions / Actual problems of organization of children's and family-san. kur.lecheniya and recreation in resort areas. Tez.nauch.-praktich.konf., Dedicated to the 100th anniversary of the resort Anan.-Anan, 1998, P.29-31 (in Russ.).
- [11] Chukov S.N. Structural and functional parameters of soil organic matter under conditions of anthropogenic impact / St. Petersburg.: St. Petersburg State University, 2011. 216 p. (in Russ.).
- [12] Adilov V.B., Berezhnov E.S., Trebukhova T.M., Trebukhov J.A. Natural curative resources Samara Region / Vopr.kurortologii, physical therapy and exercise therapy, 2008. №2. P. 45-47 (in Russ.)
- [13] Adilov V.B., Gusarev I.I. (Et al.) // On the question of the permissible content of radionuclides in mineral waters, curative mud, clay / Vopr.kurortologii, physiotherapy and LFK. 2011. №3. P. 30-33 (in Russ.)
- [14] Puntus F.A. The study of the chemical nature of humic acids of sapropel.: Avtoref.diss.kand.nauk / M., 1976, 18 p. (in Russ.)
- [15] Agapov A.I. Specific organic substances as a source of medicinal mud peloidopreparatov humic series: avtoref.dis.kand Mr. biol.nauk / -Samara, 1999.-58 P. (in Russ.)
- [16] Bachmann V.I., Ovsyannikov K.A., AD-Vadkovsky method of analysis of therapeutic mud (peloidov) / M: TsNIKIIF, 1965. 217 P. (in Russ.).
- [17] Cherepanov M.N. The group composition of organic compounds of therapeutic mud / Vopr.izucheniya medicinal mineral water, mud and klimata. M., 1976. T.31. P.142-146 (in Russ.).
- [18] Kosyanova Z.F. Chemical characterization and biological activity of humic acids of some therapeutic muds: avtoref.dis.kand.biol.nauk / M., 1985, 20 p. (in Russ.).
- [19] Krechetova E.V., Humic acid oil shale. Their properties and structure: avtoref.diss.kand.biol nauk / M., 1994, 20 p. (in Russ.).
- [20] Kan S., Tyurin A., Kurmangaliyeva Sh. Physical and chemical studies of therapeutic mud Arasan-Kundyzdy fugs // Proceedings of the NAS RK, Series of geology and engineering sciences. Almaty, Volume 5. Number 431(2018), 12-14, ISSN 2224-5278, <https://doi.org/10.32014/2018.2518-170X.3>(in Eng.).

- [21] Dospaev M. M., Bayeshov A., Zhurinov M. Z., Zhumakanova A.S., Bayeshova A.K., Syzdykova B.B., Dospaev D.M., Kakenov K.S. Physical-and-chemical regularities of forming Chrysocolla mineraling in metasilicate solution // Proceedings of the NAS RK, Series of geology and engineering sciences. Almaty, Volume 5. Number 431(2018), 107-113., ISSN 2224-5278, <https://doi.org/10.32014/2018.2518-170X.15> (in Eng.).
- [22] Bazhirov N.S., Dauletiyarov M.S., Bazhirov T.S., Serikbayev B.E., Bazhirova K.N. Research of waste of aluminium production as the raw components in technology of composite cementing materials // Proceedings of the NAS RK, Series of geology and engineering sciences. Almaty, Volume 1, Number 427 (2018), 93 – 98, ISSN 2224-5278.
- [23] Omarov B.T., Moldabekov Sh.M., Zhantasov K.T., Bolysbek A.A., Baiysbay O.P., Suigenbayeva A.Zh., Bekzhigitova K.A., Kenzhibayeva G.S., Tleuova A.B., Shapalov Sh.K. Study of the Yield of humic acids in a hydrodynamic rotary-pupulsating apparatus // Proceedings of the NAS RK, Series of geology and engineering sciences. Almaty, Volume 2, Number 428 (2018), 167 – 171. (In Eng.), ISSN 2224-5278 (in Eng.).
- [24] Zholtayev G.Zh., Zhukov N.M., Bospayev Kh.A. The theory of forecasting and evaluating the minerals and raw materials base of the Republic of // Proceedings of the NAS RK, Series of geology and engineering sciences. Almaty, Volume 2, Number 428 (2018), 193-200. ISSN 2224-5278 (in Russ.).
- [25] Katunina V.V. Ecological and biochemical activity gimatemelanovyh acids of peloids: avtoref.dis.kand.biol.nauk / Samara, 2007.-20 p. (in Russ.).
- [26] Lodygin V.D., Bezosikov V.A., Chukov S.N. Structural and functional parameters of humic substances podzonistyh bog soils podzonistyh / SPB.: Science, 2077.- P. 5-8. (in Russ.).
- [27] Popov A.I., pod.red. Ermakov E.I. Humic substances: properties, structure, education / St. Petersburg, 2004.-248p. (in Russ.).
- [28] Koponova M.M. Soil organic matter, its nature, properties, and methods of study / M.; 1963. 292 p. (in Russ.).
- [29] Kairbekov Zh., Toktamysov M.T., Zhalgasuly N., Eshova Zh.T.. Complex processing of brown coal in central Kazakhstan / Monograph. Almaty: Kazakh University, 2014. 278 p. (in Russ.).
- [30] Kairbekov Zh.K., Yermoldina E.T., Jeldybayeva I.M., Kairbekov A.Zh. Complex processing of brown coal of South Kazakhstan / Monograph: Almaty - Kazakh University, 2018. 454 b. (in Russ.).
- [31] Koponova M.M. Soil organic matter, its nature, properties and methods of study / M., 1963-292 p. (in Russ.).
- [32] Nikopova S.I., Chukov S.N. Application of the EPR method to the study of model humus-ferrous compounds / Modeling of soil-forming processes of the humid zone.-L.: Leningrad. Publishing House, 1984, P.115-128. (in Russ.)Black M.C. and McCarthy L.F. Dissolved organic matter reduces the uptake of hydrophobic organic contaminants by gills of the rainbow trout, Salmo gairdneri Environ/ Toxicol.Chem.7,p.593-600. (in Eng.).
- [33] Black M.C. and McCarthy L.F. Dissolved organic matter reduces the uptake of hydrophobic organic contaminants by gills of the rainbow trout, Salmo gairdneri Environ/ Toxicol.Chem.7,p.593-600.
- [34] Maltsev V.V. Physico-chemical properties of humic acids modified by the method of mechanical activation of caustobilot, and their interaction with biocides: Abstract. Cand. Chemical Science-Tomsk, 2010. 23 c. (in Russ.).
- [35] Avvakumova N.P., Glubokova M.N., Katunina E.E. Investigation of the antioxidant properties of peloids humic acids / Minerals.-Samara State Medical University, 2013.-C.1160-1162. (in Russ.)
- [36] Kairbekov Zh.K., Jeldybayeva I.M., Yermoldina E.T., Suymbaeva S.M. The modern view of the antioxidant activity of humic substances of peloids and their classification // Trudy IX inter. Beremzhanovsky Congress on Chemistry and Chemical Technology, Almaty. - December 9-10, 2016 - p. 315-321 (in Russ.).
- [37] Kalzhapparova I., Sabitova A., Jeldybayeva I.M. Improving the method of obtaining humic substances from peloids // International Scientific Conference of Students and Young Scientists "Ul-Farabi Elemi", Almaty, 2017, p.14 (in Russ.).
- [38] Baymold J.A., Abilmazhinova D., Jeldybayeva I.M., Kishibaev K.O. Practical application of silt sulfide mud (peloids) // Materials of the Int. Scientific-practical conference "Modern directions of development of education and science in the field of chemistry, biology, ecology and geography", Almaty, October 27, 2017, P.92-94 (in Russ.).
- [39] Abilmazhinova, DZ, Jeldybayeva, IM, Yermoldina, E.T., Kairbekov Zh.K. Method of obtaining humic substances from low-mineralized silt sulfide muds (peloids) // 4th Inter. Russian-Kazakhstan scientific practical. conference "Chemical tech. of functional materials "Almaty, April 12-13, 2018, p. 133-135 (in Russ.).

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации
в журнале смотреть на сайте:

www.nauka-nanrk.kz

<http://chemistry-technology.kz/index.php/en/arhiv>

ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)

Редакторы: *М. С. Ахметова, Т. А. Апендиев, Аленов Д. С.*
Верстка на компьютере *А. М. Кульгинбаевой*

Подписано в печать 05.06.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
5.7 п.л. Тираж 300. Заказ 3.