

**ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ГЫЛЫМ АКАДЕМИЯСЫНЫҢ

Д.В.Сокольский атындағы «Жанармай,
катализ және электрохимия институты» АҚ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
АО «Институт топлива, катализа и
электрохимии им. Д.В. Сокольского»

N E W S

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
JSC «D.V. Sokolsky institute of fuel, catalysis
and electrochemistry»

**SERIES
CHEMISTRY AND TECHNOLOGY**

6 (438)

NOVEMBER – DECEMBER 2019

PUBLISHED SINCE JANUARY 1947

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of chemistry and technologies scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Химия және технология сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруды. Web of Science зерттеушілер, авторлар, баспашилар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енүі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия химии и технологий» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Бас редакторы
х.ғ.д., проф., ҚР ҮҒА академигі **М.Ж. Жұрынов**

Редакция алқасы:

Агабеков В.Е. проф., академик (Белорус)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Ресей)
Газалиев А.М. проф., академик (Қазақстан)
Ергожин Е.Е. проф., академик (Қазақстан)
Жармағамбетова А.К. проф. (Қазақстан), бас ред. орынбасары
Жоробекова Ш.Ж. проф., академик (Қырғыстан)
Иткулова Ш.С. проф. (Қазақстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Қазақстан)
Баешов А.Б. проф., академик (Қазақстан)
Бұркітбаев М.М. проф., академик (Қазақстан)
Джусипбеков У.Ж. проф. корр.-мүшесі (Қазақстан)
Молдахметов М.З. проф., академик (Қазақстан)
Мансуров З.А. проф. (Қазақстан)
Наурызбаев М.К. проф. (Қазақстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф. академик (Қазақстан)
Стрельцов Е. проф. (Белорус)
Тәшімов Л.Т. проф., академик (Қазақстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Тәжікстан)
Фарзалиев В. проф., академик (Әзірбайжан)

«ҚР ҮҒА Хабарлары. Химия және технология сериясы».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Меншіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №1089-Ж мерзімдік басылым тіркеуіне қойылу туралы күзелік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://chemistry-technology.kz/index.php/en/archiv>

© Қазақстан Республикасының Үлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д.х.н., проф., академик НАН РК **М. Ж. Журинов**

Р е д а к ц и о н н а я к о л л е г и я:

Агабеков В.Е. проф., академик (Беларусь)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Россия)
Газалиев А.М. проф., академик (Казахстан)
Ергожин Е.Е. проф., академик (Казахстан)
Жармагамбетова А.К. проф. (Казахстан), зам. гл. ред.
Жоробекова Ш.Ж. проф., академик (Кыргызстан)
Иткулова Ш.С. проф. (Казахстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Казахстан)
Баешов А.Б. проф., академик (Казахстан)
Буркитбаев М.М. проф., академик (Казахстан)
Джусипбеков У.Ж. проф. чл.-корр. (Казахстан)
Мулдахметов М.З. проф., академик (Казахстан)
Мансуров З.А. проф. (Казахстан)
Наурызбаев М.К. проф. (Казахстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф. академик (Казахстан)
Стрельцов Е. проф. (Беларусь)
Ташимов Л.Т. проф., академик (Казахстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Таджикистан)
Фарзалиев В. проф., академик (Азербайджан)

«Известия НАН РК. Серия химии и технологии».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,

<http://chemistry-technology.kz/index.php/en/arhiv>

© Национальная академия наук Республики Казахстан, 2019

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142,
Институт органического катализа и электрохимии им. Д. В. Сокольского,
каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

NEWS**OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
SERIES CHEMISTRY AND TECHNOLOGY**

ISSN 2224-5286

<https://doi.org/10.32014/2019.2518-1491.73>

Volume 6, Number 438 (2019), 49 – 54

UDC 547.918:547.926

D. Usmanov¹, N.Sh. Ramazanov¹, U. Yusupova¹, K.Dzh. Kucherbayev^{2,3}¹Institute of Chemistry of Plant Substances AS RUz, Tashkent, Uzbekistan;²Regional Social-Innovational University, Shymkent, Kazakhstan;³JSC «South-Kazakhstan Medical Academy», Shymkent, Kazakhstan**IRIDIOIDS FROM *PHLOMIS SEVERTZOVII* AND ITS
IMMUNOSTIMULATING AND ANTITOXIC ACTIVITY**

Abstract. From the aerial parts of *Phlomis severtzovii* (*Lamiaceae*) growing in Uzbekistan, five known iridoid glycosides were isolated for the first time. On basis of UV, IR, ¹H and ¹²C NMR spectroscopy the isolated substances were identified as 6β-hydroxypolamid (1), loganin (2), pulchelloside (3), shanshiside methyl ester (4) and phlorigidoside C (5). Additionally, noticeable immunostimulating and antitoxic activity of the sum of iridoids and 6β-hydroxypolamid (1) were revealed.

Keywords: *Phlomis severtzovii*, iridoid, 6β-hydroxypolamid, loganin, pulchelloside, shanshiside methyl ester, phlorigidoside C, immunostimulating activity, antitoxic effect.

Introduction. Among the various low-molecular biologically active substances synthesized by plants, iridoids occupy a prominent place. At present time, it is clear, that this type of compounds are widespread in plant world [1]. Presence of important biological activities (antitumor activity, antimicrobial activity, etc.) of these compounds is main perspective practical reason for further investigation [2].

Iridoid glycosides have great theoretical importance as in terms of chemistry and due to their participation as precursors in alkaloid biosynthesis [3].


Therefore search of novel iridoidcontaining plant sources, development of rational scheme of isolation of these compounds, determination their chemical structure, revealing of physical, chemical parameters and useful properties of novel compounds are actual problems of the modern bioorganic chemistry [4].

The aim of our study is investigation of iridoids from *Phlomis severtzovii* (*Lamiaceae*) and their biological activity.

Materials and methods. *Phlomis severtzovii* (*Lamiaceae*) growing in Middle Asia, Tashkent and Fergana regions.

NMR spectra were acquired on a Bruker, 500 MHz, spectrometer using standard pulse sequence at ambient temperature. Chemical shifts are given in δ (ppm), and coupling constants are reported in Hz. The samples for NMR analysis were diluted in C₅D₅N.

Results and discussion. Five known iridoid glycosides were isolated from the aerial parts of *Phlomis severtzovii* collected during vegetation phase. Their chemical structure were determined on basis of analysis of UV, IR, ¹H and ¹³C NMR spectra and identified as 6β-hydroxypolamid 1, loganin 2, pulchelloside 3, shanshiside methyl ester 4 and phlorigidoside C 5 (Fig.1).



1: $R_1 = OH$, $R_2 = OH$, $R_3 = H$, $R_4 = OH$

2: $R_1 = H$, $R_2 = H$, $R_3 = OH$, $R_4 = H$

3: $R_1 = OH$, $R_2 = OH$, $R_3 = OH$, $R_4 = H$

4: $R_1 = H$, $R_2 = OH$, $R_3 = H$, $R_4 = OH$

5: $R_1 = H$, $R_2 = OH$, $R_3 + R_4 = O$

Figure 1 – Chemical structures of iridoid glycosides isolated from the aerial parts of *Phlomis severtzovii*

6 β -Hydroxypolamid (1). Compound **1** was isolated as pale yellow crystals, $C_{17}H_{26}O_{12}$, ESI-MS m/z 445.15 [$M + Na$]⁺, mp 76°C, UV (C_2H_5OH) λ_{max} 235.5 nm. IR (KBr) ν_{max} 3388 (OH), 1639 (C=O), 1551 (C=C). Yield 0.031 %.

1H NMR (Bruker 500 MHz, C_5D_5N): 6.05 (1H, s, H-1), 7.77 (1H, s, H-3), 5.55 (1H, d, H-6), 2.19 (1H, d, Ha-7), 2.55 (1H, d, Hb-7), 2.99 (1H, d, H-9), 1.44 (3H, s, H-10), 3.68 (3H, s, H-12), 5.67 (1H, d, H-1'), 4.34 (1H, t, H-2'), 4.29 (1H, t, H-3'), 4.22 (1H, t, H-4'), 4.45 (1H, ddd, H-5'), 4.37 (1H, dd, , Ha-6'), 4.28 (1H, dd, Hb-6'), 6.52 (H, br.s., OH);

^{13}C NMR (126 MHz, C_5D_5N): 93.91 (C-1), 154.92 (C-3), 115.73 (C-4), 74.4 (C-5), 72.39 (C-6), 48.58 (C-7), 73.88 (C-8), 57.59 (C-9), 26.82 (C-10), 168.64 (C-11), 52.04 (C-12), 99.01 (C-1'), 74.49 (C-2'), 78.18 (C-3'), 71.81 (C-4'), 78.67 (C-5'), 63.65 (C-6') [6].

Loganin (2). Compound **2** was isolated as white crystals. $C_{17}H_{26}O_{10}$, ESI-MS m/z 413.39 [$M + Na$]⁺, mp 222.2°C, UV (C_2H_5OH) λ_{max} 234.72 nm. IR (KBr) ν_{max} 3440 (OH), 1642 (C=O), 1446 (C=C). Yield 0.021 %.

1H NMR (Bruker 500 MHz, C_5D_5N): 5.87 (1H, d, H-1), 7.72 (1H, s, H-3), 3.59 (1H, dd, H-5), 2.07 (1H, dd, Ha-6), 2.35 (1H, dd, Hb-6), 4.04 (1H, ddd, H-7), 2.59 (1H, m, H-8), 3.13 (1H, dd, H-9), 1.16 (3H, s, H-10), 3.60 (3H, s, H-12), 5.69 (1H, d, H-1'), 4.27 (1H, t, H-2'), 4.33 (1H, t, H-3'), 4.22 (1H, t, H-4'), 4.11 (1H, ddd, H-5'), 4.19 (1H, dd, Ha-6'), 4.34 (1H, dd, Hb-6'), 6.82 (H, br.s., OH);

^{13}C NMR (126 MHz, C_5D_5N): 96.04 (C-1), 151.64 (C-3), 111.46 (C-4), 32.71 (C-5), 37.71 (C-6), 74.57 (C-7), 40.42 (C-8), 43.36 (C-9), 14.18 (C-10), 168.69 (C-11), 51.73 (C-12), 100.72 (C-1'), 75.07 (C-2'), 78.60 (C-3'), 71.81 (C-4'), 78.40 (C-5'), 63.65 (C-6') [7].

Pulchelloside (3). Compound **3** was isolated as yellow crystals. $C_{17}H_{26}O_{12}$, ESI-MS m/z 445.13 [$M + Na$]⁺, mp. 124,6°C, UV (C_2H_5OH) λ_{max} 235.76 nm. IR (KBr) ν_{max} 3437 (OH), 1640 (C=O), 1443 (C=C). Yield 0.073 %.

1H NMR (Bruker 500 MHz, C_5D_5N): 5.86 (1H, s, H-1), 7.77 (1H, s, H-3), 4.48 (1H, d, H-6), 3.72 (1H, dd, H-7), 2.41 (1H, dd, H-8), 2.90 (1H, dd, H-9), 1.22 (3H, d, H-10), 3.72 (3H, s, H-12), 5.70 (1H, d, H-1'), 4.32 (1H, dd, H-2'), 4.32 (1H, d, H-3'), 4.22 (1H, dd, H-4'), 4.46 (1H, ddd, H-5'), 4.36 (1H, dd, Ha-6'), 4.21 (1H, dd, Hb-6'), 6.57 (H, br.s., OH);

^{13}C NMR (126 MHz, C_5D_5N): 97.29 (C-1), 152.93 (C-3), 112.78 (C-4), 73.54 (C-5), 77.19 (C-6), 75.39 (C-7), 38.22 (C-8), 51.35 (C-9), 14.44 (C-10), 168.84 (C-11), 52.35 (C-12), 100.91 (C-1'), 74.94 (C-2'), 78.55 (C-3'), 71.81 (C-4'), 78.38 (C-5'), 63.65 (C-6') [8].

Shanshiside methyl ester (4). White crystals, $C_{17}H_{26}O_{11}$, ESI-MS m/z 429.38 [M + Na]⁺, mp 94.8°C, UV (C_2H_5OH) λ_{max} 236.97 nm. IR (KBr) ν_{max} 3437 (OH), 1642 (C=O), 1438 (C=C). Yield 0.013 %.

¹H NMR (Bruker 500 MHz, C_5D_5N) 6.28 (1H, s, H-1), 7.81 (1H, s, H-3), 3.33 (1H, dd, 6.48, H-5), 4.51 (1H, ddd, H-6), 2.20 (1H, dd, Ha-7), 1.80 (1H, dd, Hb-7), 2.64 (1H, dd, H-9), 1.51 (3H, s, H-10), 3.60 (3H, s, H-12), 5.77 (1H, d, H-1'), 4.30 (1H, dd, H-2'), 4.28 (1H, t, H-3'), 4.22 (1H, t, H-4'), 4.44 (1H, ddd, H-5'), 4.30 (1H, dd, Ha-6'), 4.38 (1H, dd, Hb-6'), 6.28 (1H, br.s., 6-OH);

¹³C NMR (126 MHz, C_5D_5N) 93.62 (C-1), 152.48 (C-3), 111.72 (C-4), 41.08 (C-5), 70.95 (C-6), 48.65 (C-7), 78.79 (C-8), 49.87 (C-9), 26.95 (C-10), 168.69 (C-11), 51.78 (C-12), 100.92 (C-1'), 74.81 (C-2'), 78.54 (C-3'), 71.81 (C-4'), 78.35 (C-5'), 63.65 (C-6') [9].

Phlorigidoside C (5). Colourless, $C_{17}H_{24}O_{11}$, ESI-MS m/z 427.11 [M + Na]⁺, mp 66°C, UV (C_2H_5OH) λ_{max} 238.3 nm. IR (KBr) ν_{max} 3418 (OH), 1637 (C=O), 1549 (C=C). Yield 0.028 %.

¹H NMR (Bruker 500 MHz, C_5D_5N) 6.08 (1H, d, H-1), 7.83 (1H, s, H-3), 2.81 (1H, dd, H-5), 4.17 (1H, dd, H-6), 3.48 (1H, s, H-7), 2.43 (1H, dd, H-9), 1.49 (3H, s, H-10), 3.60 (3H, s, H-12), 5.45 (1H, d, H-1'), 4.28 (1H, d, H-2'), 4.29 (1H, dd, H-3'), 4.19 (1H, dd, H-4'), 4.44 (1H, ddd, H-5'), 4.28 (1H, dd, Ha-6'), 4.38 (1H, dd, Hb-6'), 6.08 (1H, br.s., OH);

¹³C NMR (126 MHz, C_5D_5N) 93.07 (C-1), 152.55 (C-3), 106.18 (C-4), 38.20 (C-5), 73.09 (C-6), 64.23 (C-7), 65.05 (C-8), 46.80 (C-9), 16.67 (C-10), 168.47 (C-11), 51.92 (C-12), 101.46 (C-1'), 74.78 (C-2'), 78.45 (C-3'), 71.81 (C-4'), 78.35 (C-5'), 63.65 (C-6') [10].

Investigation of the biological activity. Experiments were set on white outbred mice (18-20 g). Reaction of Jerne, Nordin was used for evaluation of the preparation influence to the humoral immunity [11]. Mice were immunized with red blood cells (RBCS) of sheep and in the same day of immunization, in biological active dose the water-alcohol solutions of the iridoids sum were injected intragastric (10 мг/кг). Mice of the control group received the water-alcohol solution in the same volume. On the 5th day after immunization, the number of antibody-producing cells (APC) was determined in the spleens of mice.

Studies were performed on intact mice and mice subjected to stress. The state of stress was caused by the N. Selye "swim stress" model. To do this, mice for 45 min were placed in a tank with water, so that the water level did not allow mice to jump out.

The research results are presented in table 1.

The table shows that the SI in mice causes a pronounced ($p < 0.001$) enhancement of the process of antibody formation in response to sheep RBCS.

In the control group of immunized mice, $11,800 \pm 928.08$ APC accumulate on the spleen and 132.8 ± 2.49 APC per 1 million nucleated spleen cells (NSC). The SI increases the number of APC on the spleen to $21,400 \pm 1,369.18$, the number of APC per 1 million of NSC - to 242.0 ± 2.83 .

Table 1 – The effect of sum of iridoids from the plant *Phlomis severtzovii* on immunity indicators

Group	Control	SI
Immunized mice		
APC on spleen	11800 ± 928.08	21400 ± 1369.18
APC on 1 million of NSC	132.8 ± 2.49	242.0 ± 2.83
Immunized mice + stress		
APC on spleen	6400 ± 314.11	13200 ± 461.88
APC on 1 million of NSC	75.7 ± 3.33	151.0 ± 2.67

Under the influence of "swimming" stress, the sum of APC in mice decreases to 6400 ± 314.11 per spleen and to 75.7 ± 3.33 per 1 million NSC. The sum of iridoids increases the number of APC per spleen to $13,200 \pm 461.88$, by 1 million splenic cells - to 151.0 ± 2.67 .

Under the influence of the SI, there is a tendency to increase the mass and cellularity (total cell content) of the spleen, the mass of the thymus and lymph nodes, a significant increase in the cellularity of the thymus and lymph nodes.

As a result, it was shown that the SI obtained from the plant *Phlomis severtzovii* possesses immunostimulating activity, and is not immunotoxic.

It was previously established, that iridoids obtained from the aerial part of *Phlomis severtzovii* have hepatoprotective properties and a positive effect on the functional state of the liver [12]. Given this fact,

we investigated the effect of the SI and iridoid 6 β -hydroxypolamide on acute alcohol intoxication. Experiments were performed on white outbred mice - females weighing 20-22 g. Ethanol in the form of a 24% solution was injected intraperitoneally at a narcotic dose (4.8 g/kg). The test substances were administered orally using anatraumatic metal probe at doses of 10-25-50 mg/kg 60-70 minutes before the introduction of ethanol. Evaluation of the effectiveness of the compounds was determined by the duration of the lateral position of the animals. The control group of animals under similar conditions of experience instead of the drug was injected with sterile distilled water. The results of the research are presented in table 2.

Table 2 – The results of the influence of drugs on the narcotic action of ethanol

No	Name of preparations	dose	The duration of the lateral position in minutes	Effect in %
1	Control EtOH	4,8 g/kg	110	100%
2	SI +EtOH	10 mg/kg 4,8 g/kg	85	22,8%
3	SI + EtOH	25 mg/kg 4,8 g/kg	62,7	43%
4	SI + EtOH	50 mg/kg 4,8 g/kg	63,8	42%
5	6 β -hydroxypolamid + EtOH	10 mg/kg 4,8 g/kg	72,4	34,2%
6	6 β -hydroxypolamid + EtOH	25 mg/kg 4,8 g/kg	53,9	51%
7	6 β -hydroxypolamid + EtOH	50 mg/kg 4,8 g/kg	60,5	45%

The results of the studies (table 2) showed that in the control group of animals after applying EtOH at a dose of 4.8 g/kg in all mice came the state of anesthesia (lateral position) with an average duration of 110 minutes. Preliminary application of the SI and 6 β -hydroxypolamide at doses of 10-25-50 mg/kg shortened the state of the lateral position of the experimental mice by 22.8%-43%-42% and 34.2% -51%-45%, respectively, compared to with a control group of animals.

Consequently, the compounds studied, depending on the dose administered, have an antitoxic effect during acute alcohol intoxication.

Experimental. Air-dried and grinded plant material was extracted with MeOH three times at room temperature. After filtration, the solvent was partly removed by rotary evaporation and the extract residue was diluted with equal quantity of water. The formed precipitate in the extract was filtered. Residues of MeOH were removed by rotary evaporation. The obtained water part of the extract was extracted in consecutive order with CHCl₃ and n-BuOH. n-BuOH extract was evaporated on rotary evaporator and the n-BuOH fraction was obtained. The n-BuOH was set to column with silica gel and obtained several fractions. Re-chromatography isolation of these fractions by elution with solvent systems CHCl₃-MeOH: 1) 100÷1; 2) 50÷1; 3) 40÷1; 4) 30÷1; 5) 20÷1; 6) 15÷1; 7) 9÷1 and 8) 4÷1 yield the individual compounds: 6 β -hydroxypolamid **1**, loganin **2**, pulcheloside **3**, shanshiside methyl ester **4** and florigidoside C **5**.

In order to get novel immunomodulating preparations dried aerial parts of *Phlomis severtzovii* were extracted with EtOH at room temperature. EtOH extract was filtered and the solvent was removed on rotary evaporator. The obtained extract was diluted with water and the water mixture was treated at first with CHCl₃, then with n-BuOH five times. The solvents were removed on rotary evaporator. Yellow powder were get in yield. Enriched SI were obtained from n-BuOH fraction by treating with the solvent system CHCl₃ - EtOH 6÷1.

Conclusion. In this study, we investigated the iridoid constituent of *Phlomis severtzovii* growing in Uzbekistan and for first time we have isolated known iridoids identified on basis of UV, IR, ¹H and ¹³C

NMR spectra as 6 β -hydroxypolamid **1**, loganin **2**, pulchelloside **3**, shanshiside methyl ester **4** and phlorigidoside C **5**.

Immunostimulating activity of the sum of iridoids obtained from *Phlomis severtzovii* was revealed.

It was determined that the sum of iridoids and iridoid 6 β -hydroxypolamide have an antitoxic effect during acute alcohol intoxication.

Acknowledgments - Authors would like to thank the ICPS (Uzbek Academy of Sciences) for providing all necessary equipment, chemicals and animals for the study.

Д. Усманов¹, Н.Ш. Рамазанов¹, У. Юсупова¹, К.Дж. Кучербаев^{2,3}

¹ӨР FA Өсімдік затларының химиясы институты, Ташкент, Өзбекстан;

²Аймақтық әлеуметтік-инновациялық университет, Шымкент, Қазақстан;

³«Оңтүстік-Қазақстан медицина академиясы» АҚ, Шымкент, Қазақстан

PHLOMIS SEVERTZOVII ИРИДОИДТАРЫ ЖӘНЕ ОЛАРДЫҢ ИМУНИТЕТТИ ҮНТАЛАНДЫРУШЫ ЖӘНЕ ҰЙТТЫЛЫҚҚА ҚАРСЫ БЕЛСЕНДІЛІКТЕРІ

Аннотация. Өзбекстанда өсетін *Phlomis severtzovii* (*Lamiaceae*) жер үсті бөліктегінен бірінші рет бес иридоид гликозидтері бөліп алынды. УК, ИК, ^1H және ^{12}C ЯМР спектроскопия көмегімен олар 6 β -гидроксиполамид (1), логанин (2), пулчелозид (3), шаншизида метил эфирі (4) және флоригидозид С (5) екендігі анықталды. Қосымша иридоидтар жыйынтығы және 6 β -гидроксиполамидтың иммунитетті үнталандыруши және ұйттылыққа қарсы белсенділіктері анықталды.

Түйін сөздер: *Phlomis severtzovii*, иридоид, 6 β -гидроксиполамид, логанин, пулчелозид, шаншизида метил эфирі, флоригидозид С, иммунитетті үнталандыруши белсендік, ұйттылыққа қарсы белсенділік.

Д.Усманов¹, Н.Ш.Рамазанов¹, У.Юсупова¹, К.Дж.Кучербаев^{2,3}

¹ Институт химии растительных веществ АН РУз, Ташкент, Узбекистан;

²Региональный социально-инновационный университет, Шымкент, Казахстан;

³АО «Южно-Казахстанская медицинская академия», Шымкент, Казахстан

ИРИДОИДЫ ИЗ *PHLOMIS SEVERTZOVII* И ИХ ИММУНОСТИМУЛИРУЮЩАЯ И АНТИТОКСИЧЕСКАЯ АКТИВНОСТЬ

Аннотация. Впервые из надземных частей *Phlomis severtzovii* (*Lamiaceae*) произрастающей в Узбекистане выделены пять ранее известных иридоидных гликозидов. На основе УФ, ИК, ^1H и ^{12}C ЯМР спектроскопии выделенные вещества были идентифицированы с 6 β -гидроксиполамидом (1), логанин (2), пулчелозид (3), метиловый эфир шаншизида (4) и флуоригидозид С (5). Дополнительно, выявлены значительная иммуностимулирующая и антитоксическая активность суммы иридоидов и 6 β -гидроксиполамида (1).

Ключевые слова: *Phlomis severtzovii*, иридоид, 6 β -гидроксиполамид, логанин, пулчелозид, метиловый эфир шаншизида, флуоригидозид С, иммуностимулирующая активность, антитоксический эффект.

Information about the authors:

Durbek Usmanov - Institute of Chemistry of Plant Substances AS RUz, Tashkent, Uzbekistan, durbekusmanov@bk.ru, <https://orcid.org/0000-0002-7349-377X>;

Ugiloy Yusupova - Institute of Chemistry of Plant Substances AS RUz, Tashkent, Uzbekistan, yusupova.yu@gmail.com , <https://orcid.org/0000-0003-3401-2379>;

Nurmurod Sheralievich Ramazanov - Institute of Chemistry of Plant Substances AS RUz, Tashkent, Uzbekistan, ramazonovn@list.ru, <https://orcid.org/0000-0003-2563-626X>;

Kamal Dzhambalovich Kucherbayev - Regional Social-Innovational University, Shymkent, Kazakhstan, JSC «South-Kazakhstan Medical Academy», Shymkent, Kazakhstan, k.kamaljsci@gmail.com, <https://orcid.org/0000-0003-1981-1260>

REFERENCES

- [1] Galves M, Martin-Cordero C, Houghton PJ, Jesus Ayoso M. (2005) Antioxidant activity of methanol extracts from Plantago species. *Agricultural and food chemistry*, **53**, 1927-1933.
- [2] Yang L, Jia Z-J, Su B-N. (1997) Neolignan, phenylpropanoid and iridoid glycosides from *Pedicularis verticillata*. *Phytochemistry*, **45**, No 6, 1271-1273.
- [3] Lucjan Swigtek. Glukozydy iridoidowe w rodzinie Scrophullariaceae (1973) *Acta Polon Pharm. XXX*, No2, 203-212.
- [4] Povodish MN. (2006) Diversity of iridoids in Lamiaceae family and their biological activity. *Plant resources* [In Russian], **42**, 2, 131-149.
- [5] Maksudov MS. (1996) Iridoids from plants of Scrophulariaceae, Bignonaceae and Lamiaceae family. *Dissertation of PhD on chemistry* [In Russian], Tashkent, 127.
- [6] Stermitz FR, Abdel-Kader MS, Foderaro TA, Pomeroy M. (1994) Iridoid glycosides from some butterflies and their larval food plants. *Phytochemistry*, **37**, 4, 997-9.
- [7] Taguchi, H, Yokokawa Y, Endo T. (1973) Constituents of *Patrinia villosa*. *Yakugaku Zasshi*, **93**, 5, 607-11.
- [8] Milz S, Rimpler H. (1978) Pulchelloside I, a new iridoid from *Verbena pulchella* Sweet. *Tetrahedron Letters*, **10**, 895-8. CODEN:TELEAY ISSN:0040-4039.
- [9] Yoshio T, Hiroshi N, Hiroyuki I. (1977) Two new iridoid glucosides from *Messaenda parviflora* and *Mussaenda shikokiana*. *Phytochemistry*, **16**, 9, 1401-1404.
- [10] Kobayashi S, Mima A, Kihara M, Imakura Y. (1986) *Chem.Pharm.Bull.*, **34**, 876.
- [11] Jerne NK, Nordin AA. (1963) Plague formation in agar by single antibody producing cells. *Science*, **140**, 405.
- [12] Nabiev AN, Khushbaktova ZA, Zakhidov UV, etc. (1999) Hepatoprotective properties of iridoid glycosides in acute toxic liver damage by heliotrin in mice. *Chemical-pharm. Journ.*, **8**, 11.
- [13] A.Zh. Kaldybekova, A.T. Amangazyeva, Z.B. Halmenova, A.K. Umbetova. Development of technology for the complex isolation of biological active substances from plants of the genus *haplophyllum* A.juss., News of NAS RK. Series of Chemistry and technology. Volume 5, Number 431 (2018), 74 – 81 <https://doi.org/10.32014/2018.2518-1491.10> ISSN 2518-1491 (Online), ISSN 2224-5286 (Print).
- [14] A.A. Janibekov, T.X. Naubeev, N.Sh. Ramazanov, G.A. Sabirova, K.Dzh. Kucherbayev. Determination of chemical structure of cyclolehmanoside A from *Astragalus lemannianus*. News of NAS RK. Series of Chemistry and technology. Volume 2, Number 434 (2019), 50 – 54 <https://doi.org/10.32014/2019.2518-1491.18> ISSN 2224-5286 (Online), ISSN 2224-5286 (Print).

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации
в журнале смотреть на сайте:

www:nauka-nanrk.kz

<http://chemistry-technology.kz/index.php/en/arhiv>

ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)

Редакторы: *M. С. Ахметова, Т. А. Апендиев, Аленов Д.С.*
Верстка на компьютере *A.M. Кульгинбаевой*

Подписано в печать 05.12.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
5,25 п.л. Тираж 300. Заказ 6.

*Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*