

**NEWS**

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

**PHYSICO-MATHEMATICAL SERIES**

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.39>

Volume 4, Number 326 (2019), 14 – 21

УДК 517.9: 515.16

МРНТИ 27.31.21

**A.A. Zhadyranova**

Eurasian International Center for Theoretical Physics and Department  
of General & Theoretical Physics, Eurasian National University, Astana 010008, Kazakhstan  
a.a.zhadyranova@gmail.com

**HIERARCY OF WDVV ASSOCIATIVITY EQUATIONS**

**FOR  $n=3$  AND  $N=2$  CASE WHEN  $V_0=0$**

**WITH NEW SYSTEM  $a_t, b_t, c_t$**

**Abstract.** We investigate solutions of Witten-Dijkgraaf-E.Verlinde-H.Verlinde (WDVV) equations. The article discusses nonlinear equations of the third order for a function  $f = f(x,t)$  of two independent variables  $x,t$ . The equations of associativity reduce to the nonlinear equations of the third order for a function  $f = f(x,t)$  when prepotential  $F$  dependet of the metric  $\eta$ . In this work we consider the WDVV equation for  $n = 3$  case with an antidiagonal metric  $\eta$ . The solution of some cases of hierarchy equations of associativity illustrated. Lax pairs for the system of three equations, that contains the equation of associativity are written to find the hierarchy of associativity equation. Using the compatibility condition are found the relations between the matrices  $U, V_2, V_1$ . The elements of matrix  $V_2$  are found with the expression of  $z_{ij}$  and independent and dependent variables for the matrix  $V_2$ . Also solving elements of matrix  $V_1$  expressed through  $y_{ij}$  and independent and dependent variables for the matrix  $V_1$ . We accepted that elements of matrix  $V_0$  are zero. In the physical setting the solutions of WDVV describe moduli space of topological conformal field theories [1, 2]. Let us introduce new variables  $a, b, c$ . In the above variables the nonlinear equations of the third order for a function  $f = f(x,t)$  we rewritten as a new system of three equations. Expressed are variables  $a_t, b_t, c_t$  of three equations are written with the help of matrix elements  $z_{ij}, y_{ij}$ .

**Key words:** equations of Witten-Dijkgraaf-E.Verlinde-H.Verlinde, the equations of associativity, nonlinear equations of the third order, antidiagonal metric, the Lax pair, the compatibility condition, independent elements, dependent variables, system with equations.

**Introduction.** The WDVV equations, in general, have the following form [3, 4, 5]:

$$\frac{\partial^3 F}{\partial t^i \partial t^j \partial t^p} \eta^{pq} \frac{\partial^3 F}{\partial t^q \partial t^k \partial t^r} = \frac{\partial^3 F}{\partial t^j \partial t^k \partial t^p} \eta^{pq} \frac{\partial^3 F}{\partial t^i \partial t^q \partial t^r}, \quad \forall i, j, k, r \in \{1, \dots, n\},$$

where  $F$  is a prepotential,  $\eta$  is a metric. The coordinates  $t^i$  can be linearly rearranged so that the metric,  $\eta$ , is antidiagonal [6], i.e.

$$\eta = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

In this work we consider the WDVV equation for  $n = 3$  case with an antidiagonal metric  $\eta$  [7]. In this case, two types of dependence of the function  $F$  on the fixed variable  $t^1$  were found by Dubrovin [8, 9, 10] which are

$$F = \frac{1}{2}(t^1)^2 t^3 + \frac{1}{2}t^1(t^2)^2 + f(t^2, t^3) \quad (1)$$

and

$$F = \frac{1}{6}(t^1)^3 + t^1 t^2 t^3 + f(t^2, t^3).$$

For these cases the equations of associativity reduce to the following two nonlinear equations of the third order for a function  $f = f(x, t)$  of two independent variables ( $x = t^2, t = t^3$ ):

$$f_{tt} = f_{xxt}^2 - f_{xxx} f_{xtt} \quad (2)$$

and

$$f_{xxx} f_{ttt} - f_{xxt} f_{xtt} = 1,$$

correspondingly.

The function  $F$  in equation (1) has the form from the law of multiplication in the three-dimensional algebra  $A_t$  with the basis  $e_1 = 1, e_2, e_3$  [3]. Every basis is a complete uniformly minimal system [11].

In this work, we consider the solution (1). Let us introduce new variables  $a, b, c$  as follows [12, 13]:

$$a = f_{xxx}, \quad b = f_{xxt}, \quad c = f_{xtt}.$$

In the above variables the equation (2) can be rewritten as a system of three equations as follows:

$$\begin{aligned} a_t &= b_x, \\ b_t &= c_x, \\ c_t &= (b^2 - ac)_x. \end{aligned} \quad (3)$$

The Lax pair for the system (3) is given by [8]

$$\begin{aligned} \Psi_x &= \lambda U \Psi, \\ \Psi_t &= \lambda V \Psi, \end{aligned} \quad (4)$$

where  $U$  is given by

$$U = \begin{pmatrix} 0 & 1 & 0 \\ b & a & 1 \\ c & b & 0 \end{pmatrix}$$

and  $V$  is given by

$$V = \begin{pmatrix} 0 & 0 & 1 \\ c & b & 0 \\ (b^2 - ac) & c & 0 \end{pmatrix}.$$

The compatibility condition for the system (4) is given by

$$\begin{aligned} U_t &= V_x, \\ [U, V] &= 0. \end{aligned}$$

In the following sections we work with the new system (3).

**Methods.** The solution to a hierarchy for  $N = 1$  case corresponds to the system of equations (3). Hierarchy for  $N = 2$  case when  $V_0 \neq 0$  is given in the work [14]

In this section we consider a hierarchy for  $N = 2$  case when  $V_0 = 0$  and the following system

$$\begin{aligned} a_t &= \varepsilon_1 b_x + \varepsilon_2 F_x, \\ b_t &= \varepsilon_1 c_x + \varepsilon_2 H_x, \\ c_t &= \varepsilon_1 (b^2 - ac)_x + \varepsilon_2 G_x. \end{aligned} \quad (5)$$

The Lax representation of the above system is same as before in the work [13].

In particular, for  $N = 2$  case when  $V_0 = 0$  we have

$$\begin{aligned} \Psi_x &= \lambda U \Psi, \\ \Psi_t &= (\lambda^2 V_2 + \lambda V_1) \Psi = V \Psi \end{aligned}$$

The compatibility condition of (4) is given by

$$\lambda U_t - V_x + \lambda [U, V] = 0.$$

The compatibility condition of the Lax representation is given by the system

$$[U, V_2] = 0, \quad (6)$$

$$U_t = V_{1x}, \quad (7)$$

$$V_{2x} = [U, V_1] \quad (8)$$

**Statement of problem.** We first consider the second equation of the system and let  $V_1$  to be given by

$$V_1 = \begin{pmatrix} y_{11} & y_{12} & y_{13} \\ y_{21} & y_{22} & y_{23} \\ y_{31} & y_{32} & y_{33} \end{pmatrix}.$$

From the above system it follows that  $y_{11}, y_{12}, y_{13}, y_{21}, y_{22}, y_{23}, y_{31}, y_{32}, y_{33}$  are constants w.r.t.  $x$ . Writing a system with equations for  $a_t, b_t, c_t$  only yields

$$\begin{aligned} a_t &= y_{22x}, \\ b_t &= y_{21x}, \\ b_t &= y_{32x}, \\ c_t &= y_{31x}. \end{aligned} \quad (9)$$

Now we equate similar terms in the systems (5) and (9), i.e. we have a system

$$\begin{aligned} a_t &= y_{22x} = \varepsilon_1 b_x + \varepsilon_2 F_x, \\ b_t &= y_{21x} = \varepsilon_1 c_x + \varepsilon_2 H_x, \\ b_t &= y_{32x} = \varepsilon_1 c_x + \varepsilon_2 H_x, \\ c_t &= y_{31x} = \varepsilon_1 (b^2 - ac)_x + \varepsilon_2 G_x. \end{aligned} \quad (10)$$

**Scheme of the method and reduction to equivalent problem.** From the above system (10) we find the following

$$\begin{aligned} y_{22} &= \varepsilon_1 b + \varepsilon_2 F, \\ y_{21} &= \varepsilon_1 c + \varepsilon_2 H, \\ y_{32} &= \varepsilon_1 c + \varepsilon_2 H, \\ y_{31} &= \varepsilon_1 (b^2 - ac) + \varepsilon_2 G. \end{aligned}$$

Thus the matrix  $V_1$  has the form

$$V_1 = \begin{pmatrix} y_{11} & y_{12} & y_{13} \\ \varepsilon_1 c + \varepsilon_2 H & \varepsilon_1 b + \varepsilon_2 F & y_{23} \\ \varepsilon_1 (b^2 - ac) + \varepsilon_2 G & \varepsilon_1 c + \varepsilon_2 H & y_{33} \end{pmatrix}. \quad (11)$$

Now we solve the equation (6). Denote  $V_2$  as follows:

$$V_2 = \begin{pmatrix} z_{11} & z_{12} & z_{13} \\ z_{21} & z_{22} & z_{23} \\ z_{31} & z_{32} & z_{33} \end{pmatrix},$$

Plugging  $U$ ,  $V_2$  into (6) we obtain the following relations:

$$\begin{aligned} z_{23} &= z_{12}, \\ z_{32} &= z_{21}, \\ z_{33} &= z_{11}. \end{aligned}$$

Hence, we are left with the equations

$$\begin{aligned} z_{21} &= bz_{12} + cz_{13}, \\ z_{22} &= z_{11} + az_{12} + bz_{13}, \\ z_{31} &= cz_{12} + (b^2 - ac)z_{13}. \end{aligned}$$

Thus the matrix  $V_2$  has the form

$$V_2 = \begin{pmatrix} z_{11} & z_{12} & z_{13} \\ bz_{12} + cz_{13} & z_{11} + az_{12} + bz_{13} & z_{12} \\ cz_{12} + (b^2 - ac)z_{13} & bz_{12} + cz_{13} & z_{11} \end{pmatrix}.$$

Hence, only  $z_{11}, z_{12}, z_{13}$  are independent elements of  $V_2$ , and the other elements can be written in terms of them.

Now let us find the elements of  $V_1$  in (11). To do so we use the equation (8). First we evaluate  $[U, V_1]$ .

We have elementwise yields the following system:

- 11:  $z_{11x} = \varepsilon_1 c + \varepsilon_2 H - b y_{12} - c y_{13}$ ,
- 12:  $z_{12x} = \varepsilon_1 b + \varepsilon_2 F - y_{11} - a y_{12} - b y_{13}$ ,
- 13:  $z_{13x} = y_{23} - y_{12}$ ,
- 21:  $b_x z_{12} + b z_{12x} + c_x z_{13} + c z_{13x} = b y_{11} + a(\varepsilon_1 c + \varepsilon_2 H) + (\varepsilon_1(b^2 - ac) + \varepsilon_2 G) - b(\varepsilon_1 b + \varepsilon_2 F) - c y_{23}$ ,
- 22:  $z_{11x} + a_x z_{12} + a z_{12x} + b_x z_{13} + b z_{13x} = b y_{12} - b y_{23}$ ,
- 23:  $z_{12x} = b y_{13} + a y_{23} + y_{33} - (\varepsilon_1 b + \varepsilon_2 F)$ ,
- 31:  $c_x z_{12} + c z_{12x} + (b^2 - ac)_x z_{13} + (b^2 - ac) z_{13x} = c y_{11} - c y_{33}$ ,
- 32:  $b_x z_{12} + b z_{12x} + c_x z_{13} + c z_{13x} = c y_{12} + b(\varepsilon_1 b + \varepsilon_2 F) - (\varepsilon_1(b^2 - ac) + \varepsilon_2 G) - a(\varepsilon_1 c + \varepsilon_2 H) - b y_{33}$ ,
- 33:  $z_{11x} = c y_{13} + b y_{23} - (\varepsilon_1 c + \varepsilon_2 H)$ .

Now let us express  $\varepsilon_1 c + \varepsilon_2 H$ ,  $\varepsilon_1 b + \varepsilon_2 F$ ,  $y_{23}$  in the element 11, 12, 13 of the above system.

$$\begin{aligned}\varepsilon_1 c + \varepsilon_2 H &= z_{11x} + b y_{12} + c y_{13}, \\ \varepsilon_1 b + \varepsilon_2 F &= z_{12x} + y_{11} + a y_{12} + b y_{13}, \\ y_{23} &= z_{13x} + y_{12}.\end{aligned}$$

Now let us express  $\varepsilon_1(b^2 - ac) + \varepsilon_2 G$  in the element 21 and substitute the values for  $\varepsilon_1 c + \varepsilon_2 H$ ,  $\varepsilon_1 b + \varepsilon_2 F$ ,  $y_{23}$

$$\varepsilon_1(b^2 - ac) + \varepsilon_2 G = b_x z_{12} + b z_{12x} + c_x z_{13} + 2 c z_{13x} - a z_{11x} + b z_{12x} + (b^2 - ac) y_{13} + c y_{12}$$

Now let us express  $y_{33}$  in the element 23 and substitute the values for  $\varepsilon_1 b + \varepsilon_2 F$ ,  $y_{23}$

$$y_{33} = 2 z_{12x} - a z_{13x} + y_{11}$$

Hence, dependent elements of  $V_1$  are given by:

$$\begin{aligned}\varepsilon_1(b^2 - ac) + \varepsilon_2 G &= b_x z_{12} + b z_{12x} + c_x z_{13} + 2 c z_{13x} - a z_{11x} + b z_{12x} + (b^2 - ac) y_{13} + c y_{12}, \\ \varepsilon_1 c + \varepsilon_2 H &= z_{11x} + b y_{12} + c y_{13}, \\ \varepsilon_1 b + \varepsilon_2 F &= z_{12x} + y_{11} + a y_{12} + b y_{13}, \\ y_{23} &= z_{13x} + y_{12}, \\ y_{33} &= 2 z_{12x} - a z_{13x} + y_{11}.\end{aligned}\tag{12}$$

Now let us rewrite the element 22 by substituting the values for  $y_{23}$ . So we have

$$z_{11x} + 2 b z_{13x} + a_x z_{12} + a z_{12x} + b_x z_{13} = 0$$

Now let us rewrite the element 31 by substituting the values for  $y_{33}$ . So we have

$$c_x z_{12} + 3 c z_{12x} + (b^2 - ac)_x z_{13} + (b^2 - 2ac) z_{13x} = 0$$

Now let us rewrite the element 32 by substituting the values for  $\varepsilon_1 b + \varepsilon_2 F$ ,  $\varepsilon_1(b^2 - ac) + \varepsilon_2 G$ ,  $\varepsilon_1 c + \varepsilon_2 H$ ,  $y_{33}$ . So we have

$$2 b_x z_{12} + 4 b z_{12x} + 2 c_x z_{13} + (3c - ab) z_{13x} = 0$$

Now let us rewrite the element 33 by substituting the values for  $y_{23}$ ,  $\varepsilon_1 c + \varepsilon_2 H$

$$2z_{11x} - bz_{13x} = 0$$

Also, the independent variables  $z_{11}, z_{12}, z_{13}$  of the matrix  $V_2$  have to satisfy the following system of equations:

$$\begin{aligned} z_{11x} + 2bz_{13x} + a_x z_{12} + az_{12x} + b_x z_{13} &= 0, \\ c_x z_{12} + 3cz_{12x} + (b^2 - ac)_x z_{13} + (b^2 - 2ac)z_{13x} &= 0, \\ 2b_x z_{12} + 4bz_{12x} + 2c_x z_{13} + (3c - ab)z_{13x} &= 0, \\ 2z_{11x} - bz_{13x} &= 0. \end{aligned} \quad (13)$$

From the above system (13) it follows that

$$z_{13x} = \left( \frac{4a_x b - 2ab_x}{3ac - a^2b - 10b^2} \right) z_{12} + \left( \frac{4bb_x - 2ac_x}{3ac - a^2b - 10b^2} \right) z_{13} \quad (14)$$

$$z_{12x} = \left( -\frac{c_x}{3c} - \frac{b^2 - 2ac}{3c} \cdot \frac{4a_x b - 2ab_x}{3ac - a^2b - 10b^2} \right) z_{12} + \left( -\frac{b^2 - 2ac}{3c} \cdot \frac{4bb_x - 2ac_x}{3ac - a^2b - 10b^2} - \frac{(b^2 - ac)_x}{3c} \right) z_{13} \quad (15)$$

**Results.** Using necessary terms in the system (12) in (10), we obtain

$$\begin{aligned} a_t &= \frac{a_x z_{13x}}{2} + a_x y_{12} + b_x y_{13}, \\ b_t &= \frac{b_x z_{13x}}{2} + b_x y_{12} + c_x y_{13}, \\ c_t &= b_{xx} z_{12} + 3b_x z_{12x} + c_{xx} z_{13} + (a_x b + 3c_x - \frac{ab_x}{2}) z_{13x} - a_x z_{11x} + (b^2 - ac)_x y_{13} + c_x y_{12} \end{aligned} \quad (16)$$

We plug  $z_{11x}, z_{12x}, z_{13x}$  in (13), (15), (14) into (16) and obtain the following equation

$$\begin{aligned} a_t &= \left( \frac{2ba_x^2 - aa_x b_x}{3ac - a^2b - 10b^2} \right) z_{12} + \left( \frac{2ba_x b_x - aa_x c_x}{3ac - a^2b - 10b^2} \right) z_{13} + a_x y_{12} + b_x y_{13}, \\ b_t &= \left( \frac{2ba_x b_x - ab_x^2}{3ac - a^2b - 10b^2} \right) z_{12} + \left( \frac{2bb_x^2 - ab_x c_x}{3ac - a^2b - 10b^2} \right) z_{13} + b_x y_{12} + c_x y_{13}, \\ c_t &= \left( b_{xx} - \frac{b_x c_x}{c} + \frac{5abcq_x b_x - 4b^3 a_x b_x + 2ab^2 b_x^2 - 3a^2 cb_x^2 + 2b^2 ca_x^2 + 12bcq_x c_x - 6acb_x c_x}{3ac^2 - a^2 bc - 10b^2 c} \right) z_{12} \\ &+ \left( c_{xx} - \frac{b_x(b^2 - ac)_x}{c} + \frac{6abch_x^2 - 4b^3 b_x^2 + 2ab^2 b_x c_x - 3a^2 cb_x c_x + 2b^2 ca_x b_x + 12bcq_x c_x - abcq_x c_x - 6acc_x^2}{3ac^2 - a^2 bc - 10b^2 c} \right) z_{13} \\ &+ (b^2 - ac)_x y_{13} + c_x y_{12} \end{aligned} \quad (17)$$

**Conclusion.** The solution to a hierarchy for  $N = 2$  case when system is given by (5) corresponds to the system of equations (17).

So, we considered of some cases of hierarchy of WDVV associativity equations. Lax pairs for the system of three equations, that contained the equation of associativity written to find the hierarchy of associativity equation. Using the compatibility condition are found the relations between the matrices  $U$ ,  $V_2$ ,  $V_1$ . Thus, we obtained the elements of the matrices  $V_2$ ,  $V_1$  for case  $N = 2$  when  $V_0=0$  and the above system  $a_t, b_t, c_t$ . It was found, that only  $z_{11}, z_{12}, z_{13}$  are independent elements of  $V_2$ , and the other elements can be written in terms of them. From the above system it follows that  $y_{11}, y_{12}, y_{13}, y_{23}, y_{33}$

are constants w.r.t.  $\mathcal{X}$ . It is found, that  $y_{11}, y_{12}, y_{13}$  are independent elements of  $V_1$ , and the other elements can be written in terms of them and  $z_{11}, z_{12}, z_{13}$ . Expressed are variables  $a_t, b_t, c_t$  of three equations are written with the help of matrix elements  $z_{ij}, y_{ij}$ .

**Acknowledgments.** I express gratitude to Professor R. Myrzakulov for useful discussions and advices. The work is performed under the financial support of the scientific and technical program BR05236277 "Investigation of some problems of astrophysics and cosmology in the framework of the Einstein and non-Einstein theories of gravity", 2018.

УДК 517.9: 515.16  
МРНТИ 27.31.21

**А.А. Жадыранова<sup>1</sup>**

<sup>1</sup>Л.Н.Гумилев атындағы Еуразия ұлттық университетінің жалпы  
және теориялық физика кафедрасы, Астана, Қазақстан

## **n = 3 және N = 2 жағдайлары үшін енгізгілген жаңа жүйе $a_t, b_t, c_t$ V<sub>0</sub> = 0** **БОЛҒАНДАҒЫ WDVV АССОЦИАТИВІЛІК ТЕНДЕУІНІҢ ИЕРАРХИЯСЫ**

**Аннотация.** Берілген мақалада Виттен – Диджграф - Е.Верлинде - Г.Верлинде (ВДВВ) тендеулері зерттеледі. Бұл жұмыста x, t тәуелсіз айнымалыларынан тұратын  $f = f(x,t)$  функциясы үшін үшінші ретті сзыбықты емес тендеулер талқыланады. Тәуелсіз x, t айнымалыларынан тұратын  $f = f(x,t)$  функциясы үшін үшінші ретті сзыбықты емес тендеулер F потенциалы  $\eta$  метрикасымен байланысты болғанда келтіріледі. Сонымен қатар ассоциативтілік тендеулер иерархиясының бірнеше шешімдері сипатталады. Ассоциативтілік тендеулерінің иерархиясын табу мақсатында ассоциативтілік тендеулерінен құралған тендеулер жүйесі үшін Лакс жұптары жазылды. Сәйкестік шартының қолдану арқылы  $U, V_2, V_1$  матрицалары арасындағы қатынастар анықталды.  $z_{ij}$  арқылы өрнектелген  $V_2$  матрицасының элементтері мен  $V_2$  матрицасының тәуелді және тәуелсіз айнымалылары есептелінді.  $u_{ij}$  арқылы өрнектелген  $V_1$  матрицасының элементтері мен  $V_1$  матрицасының тәуелді және тәуелсіз айнымалылары табылды. Сонымен қатар  $V_0$  матрицасының элементтері нөлге тең деп алынды. Физикалық қолданылуда WDVV ассоциативтілік тендеуінің шешімі өрістің топологиялық конформдық теориясының модульдерінің кеңістігін сипаттайтыны. Жаңа айнымалылар енгізілген. Жаңа айнымалыларда  $f=f(x,t)$  функциясы үшін үшінші ретті сзыбықты емес тендеулер жаңа жүйе арқылы жазылған. Тендеулер жүйесінен тұратын  $a_t, b_t, c_t$  айнымалылары  $z_{ij}, u_{ij}$  матрицалық элементтері арқылы өрнектеліп жазылды.

**Түйін сөздер:** Виттен-Диджграф-Е.Верлинде-Г.Верлинде тендеулері, ассоциативтілік тендеуі, үшінші ретті сзыбықты емес тендеулер, антидиагональ метрика, Лакс жұптары, үйлесімділік шарты, тәуелсіз элементтер, тәуелді айнымалылар, тендеулер жүйесі.

УДК 517.9: 515.16  
МРНТИ 27.31.21

**А.А. Жадыранова<sup>1</sup>**

<sup>1</sup>Кафедра общей и теоретической физики Евразийского национального университета имени Л.Н.Гумилева,  
Астана, Казахстан

## **ИЕРАРХИЯ УРАВНЕНИЙ АССОЦИАТИВНОСТИ WDVV ДЛЯ СЛУЧАЯ n = 3 И N = 2 ПРИ V<sub>0</sub> = 0 С НОВОЙ СИСТЕМОЙ $a_t, b_t, c_t$**

**Аннотация.** В данной статье исследуются уравнения Виттена-Диджграфа-Е.Верлинде-Г.Верлинде (ВДВВ). В работе обсуждаются нелинейные уравнения третьего порядка для функции  $f = f(x,t)$  двух независимых переменных x,t. Уравнения ассоциативности сводятся к нелинейным уравнениям третьего порядка для функции  $f = f(x,t)$  когда потенциал функции F связан с метрикой  $\eta$ . В этой работе рассматривается уравнение WDVV для случая n = 3 с антидиагональной метрикой  $\eta$ . Описано решение некоторых случаев иерархии уравнений ассоциативности. Для нахождения иерархии уравнений ассоциативности были записаны пары Лакса для системы из трех уравнений, которая содержит уравнения

ассоциативности. С применением условия совместности найдены соотношения между матрицами  $U, V_2, V_1$ . Были вычислены элементы матрицы  $V_2$ , выраженные через  $z_{ij}$ , независимые и зависимые переменные матрицы  $V_2$ . Также были найдены элементы матрицы  $V_1$ , выраженные через  $u_{ij}$ , независимые и зависимые переменные матрицы  $V_1$ . Элементы матрицы  $V_0$  равны 0. В физическом приложении решение уравнения ассоциативности WDVV описывает пространство модулей топологических конформных теорий поля. Введены новые переменные  $a, b, c$ . В новых переменных нелинейные уравнения третьего порядка для функции  $f = f(x,t)$  записаны через новую систему трёх уравнений. Выраженные переменные  $a_t, b_t, c_t$  системы из трех уравнений были записаны через матричные элементы  $z_{ij}, u_{ij}$ .

**Ключевые слова:** уравнения Виттена-Дижграфа-Е.Верлинде-Г.Верлинде, уравнения ассоциативности, нелинейные уравнения третьего порядка, антидиагональная метрика, пары Лакса, условие совместности, независимые элементы, зависимые переменные, система с уравнениями.

#### Information about authors:

Zhadyranova A.A. - PhD student of the department of general and theoretical physics, L.N. Gumilyov Eurasian National University, Satpayev str., Astana, Kazakhstan. E-mail: a.a.zhadyranova@gmail.com

#### REFERENCES

- [1] Kontsevich M., Manin Yu., Gromov-Witten classes, quantum cohomology and enumerative geometry, Mirror symmetry, II, 607-653, AMS/IP Stud. Adv. Math., 1, Amer. Math. Soc., Providence, RI, (1997).
- [2] Kontsevich M., Manin Yu., Relations between the correlators of the topological sigma-model coupled to gravity, Comm. Math. Phys., 196 (1998), no. 2, 385-398.
- [3] Dubrovin B.A. Geometry of 2D topological field theories, Springer Lecture Notes in Math. 1620, 120-348, 1996. [arXiv:hep-th/9407018]
- [4] Dijkgraaf R., Verlinde E., Verlinde H., Notes on topological string theory and 2D quantum gravity, Nucl. Phys. B 352 (1991) 59.
- [5] Witten E., On the structure of the topological phase of two-dimensional gravity, Nucl. Phys.B 340 (1990) 281-332.
- [6] Hertling C., Frobenius manifolds and moduli spaces for singularities, Cambridge University Press, Cambridge (UK), 2002.
- [7] Hertling C., Manin Y., Weak Frobenius manifolds, Internat. Math. Res. Notices, no. 6 (1999), 277-286.
- [8] Mokhov O.I., Ferapontov Y.V. Equations of Associativity in Two-Dimensional Topological Field Theory as Integrable Hamiltonian Nondiagonalizable Systems of Hydrodynamic Type, Functional analysis and its applications 30(3), 1995. [arXiv:hep-th/9505180]
- [9] Dubrovin B.A., Novikov S.P., Hydrodynamics of weakly deformed soliton lattices. Differential geometry and Hamiltonian theory, Uspekhi Mat.Nauk. 44 (1989), 29-98. English translation in Russ. Math. Surveys 44 (1989), 35-124.
- [10] Dubrovin B.A., On almost duality for Frobenius manifolds, Amer. Math. Soc. Transl. 212 (2004) 75-132.
- [11] A.Sh. Shaldanbayev , A.A. Shaldanbayeva , B.A. Shaldanbay (2019) On projectional orthogonal basis of a linear non-self -adjoint operator. News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series. Volume 2, Number 324 (2019), PP, 79–89. <https://doi.org/10.32014/2019.2518-1726.15> ISSN 2518-1726 (Online), ISSN 1991-346X (Print).
- [12] Mokhov O.I. Symplectic and poisson geometry on loop spaces of manifolds and nonlinear equations, Translations of the American Mathematical Society-Series 2 170, 121-152, 1995. [arXiv:hep-th/9503076]
- [13] Ferapontov E.V., Mokhov O.I., Nonlocal Hamiltonian operators of hydrodynamic type that are connected with metrics of constant curvature, Russ. Math. Surv. 45 (1990), no. 3, 218-219.
- [14] A.A. Zhadyranova, Zh.R. Myrzakul, Y.Ye. Anuarbekova Hierarchy of WDVV associativity equations for  $n = 3$  case and  $N = 2$  when  $V_0 \neq 0$  [Ierarchiya WDVV uravneniya dlya  $n = 3$  i  $N = 2$  sluchaya, kogda  $V_0 \neq 0$ ] Bulletin of L.N. Gumilyov Eurasian National University [Vestnik Evrazijskogo naciona'l'nogo universiteta imeni L.N. Gumileva]. 4(125), 60-66, 2018.